Rocky Mountain Medical Journal

Directory

Number



Vol. 49—No. 2 Part 1

\$3.50 Per Year

35c Per Copy

CRANIOCEREBRAL TRAUMA (FEB 13)

MEDICINE'S OPPORTUNITY OF A LIFETIME — CONCENITAL ARTRESIA

POSTOPERATIVE PULMONARY INFECTIONS
KERATITIS



". . . and send it right away"

When immediate therapy is vital to the patient, critical moments are saved if your pharmacy has your first prescription choice on hand; in addition, your own valuable time is not wasted by unnecessary telephone calls and second selections.

Throughout the country, Lilly products are most widely stocked, most readily available. Save time; make Lilly your first choice.

Lilly

ELI LILLY AND COMPANY . INDIANAPOLIS 6, INDIANA, U. S. A.

in thema

DILANTIN®

"...little depression

...strong opposition to major convulsions."

PARKE, DAVIS & COMPANY

nemanagement of epilepsy

"The introduction of diphenylhydantoin was a marked advance in therapy, because this drug, although distantly related to the barbiturates, produced little depression while exerting strong opposition to major convulsions."*

Extensive clinical experience confirms the finding that DILANTIN—producing little or no depression—prevents seizures or greatly reduces their number and severity in the majority of epileptic patients.

DILANTIN Sodium (diphenylhydantoin sodium, Parke-Davis) is available in Kapseals® of 0.03 Gm. (½ gr.) and 0.1 Gm. (½ gr.) in bottles of 100 and 1000.

*Cutting, W. C.: A Manual of Clinical Therapeutics, ed. 2, Philadelphia, W. B. Saunders & Co.; 1948, p. 484.








Table of Contents

VOLUME 49

NUMBER 2

FEBRUARY, 1952

TEDROAKI, 1732	
Editorials Pa	age
Russians Attend a Medical Meeting	111
Report of Delegates to the A.M.A	
Report of Delegates to the A.M.A	112
*	
Original Articles	
Craniocerebral Trauma, Alexander C. Johnson, M.D.	113
Medicine's Opportunity of a Lifetime, Ervin A. Hinds, M.D.	120
Prevention of Postoperative Pulmonary Infections, John A. Dixon, M.D., and Earle B. Mahoney, M.D.	122
Congenital Atresia of the Bile Ducts, Paul D. Keller, M.D.	126
*	
Case Report New Drugs in the Treatment of Keratitis	129
*	
Organization	
This Is Late but Too Good to Miss	130
Colorado	
Component Societies	
Obituaries	
Wyoming News Notes	134
New Mexico Obituaries	
Blue Cross and Blue Shield.	
Colorado State Health Department	
Colorado Medical School Notes	136
Book Corner	144



PRIODAX (brand of Iodoalphionic Acid U.S.P.) CHOLECYSTOGRAPHY

presents

gallbladder shadows of great clarity free from confusing shadows

assisted by

a simple preparatory routine and tolerance of the highest degree.



schering corporation, BLOOMFIELD, N. J.



ROCKY MOUNTAIN MEDICAL JOURNAL

Title Registered, U. S. Patent Office

Publication Office: 835 Republic Building (1612 Tremont Place), Denver 2, Colorado Telephone AComa 0547

EDITORIAL BOARD

Colorado: Douglas W. Macomber, M.D., Scientific Editor, 1800 High St., Denver; Lyman W. Mason, M.D., Associate Editor, 1214 Republic Bldg., Denver (Chairman of Editorial Board).

Montana: Raymond F. Peterson, M.D., Scientific Editor, 9 W. Granite St., Butte; L. Russell Hegland, Associate Editor, 240 Stapleton Bldg., Billings.

New Mexico: Carl H. Gellenthien, M.D., Scientific Editor, Valmora, New Mexico; Ralph R. Marshall, Associate Editor, 323 First National Bank Bldg., Albuquerque.

Utah: Richard P. Middleton, M.D., Scientific Editor, Boston Bldg., Salt Lake City; W. H. Tibbals, Associate Editor, 42 South Fifth East St., Salt Lake City.

Wyoming: Franklin D. Yoder, M.D., Scientific Editor, State Capitol Building, Cheyenne; Arthur R. Abbey, Associate Editor, P. O. Box 897, Cheyenne.

Managing Editor: Harvey T. Sethman, 835 Republic Bldg., Denver. Business Manager: Helen Kearney, 835 Republic Bldg., Denver.

Business Manager: Helen Kearney, 835 Republic
Ownership and Sponsorship: The Rocky Mountain
Medical Journal is owned by the Colorado State
Medical Society and is published monthly as a nonprofit enterprise for the mutual benefit of the organizations which jointly sponsor it. It is published
under the direction of the Board of Trustees of the
Colorado State Medical Society, assisted by an Editorial Board representing the sponsoring organizations. It is the Official Journal of the Colorado State
Medical Society, the Montana Medical Association,
the New Mexico Medical Society, the Utah State
Medical Association, the Wyoming State Medical
Society, the Rocky Mountain Medical Conference,
and the Colorado Hospital Association.

Manuscripts: Scientific Articles, Case Reports, etc.,
from any state for which this is the Official Journal
should be submitted to the Scientific Editor for that
state as mamed in the Editorial Board, above. Other
material from any participating state should be submitted to the Associate Editor for that state as
named above. Manuscripts from outside the Rocky
Mountain area should be sent direct to the Journal
office. Manuscripts must be typewritten, double or
triple spaced, using only one side of each sheet. It

Advertising: National representatives: The State Journal Advertising Bureau, 535 North Dearborn Street, Chicago 10, Ill. Local advertising from firms in the Rocky Mountain area should be submitted to the Associate Editor of the appropriate state or to the Journal office. Advertising forms close on the 20th of the month preceding publication; allow ten days additional to insure submitting proofs for approval.

Subscription: \$3.50 per year in advance, postpaid in the United States and its possessions; single copy, 35c plus postage. Subscription is included in medical society dues of sponsoring state medical

Copyright: This Journal is copyright, 1952, by the Colorado State Medical Society. Requests for permission to reproduce anything from the columns of this Journal should be addressed to the Journal office.

Second Class Matter: Entered as second class matter Jan. 22, 1906, at the Post Office at Denver, Colounder the Act of Congress of March 3, 1879. Accepted for mailing at special rates of postage provided for in Section 1103, Act. of Oct. 3, 1917; authorized July 17, 1918.

Collection

of your

All reports show a trend toward slower and harder collections in the

At the first sign of neglect you will save money if they are turned over to us for collection.

Comparison of collection results, backed by 35 years of experience, proves you obtain greater results at less cost, when you list your accounts


with

The American Medical and Dental Association

2106 Broadway

TAbor 2331

Denver, Colorado



well-trained" amine the

> Seven years of search and synthesis in the Upjohn laboratories culminated in the development of Orthoxine Hydrochloride-a bronchodilator and antispasmodic. Orthoxine's dependable action gives relief to the asthma patient with minimal vasopressor and psychomotor stimulation.

Orthoxine Hydrochloride is administered orally.

Orthoxine

Bottles of 100 and 500 tablets.

Orthoxine Hydrochloride (100 mg.) Tablets contain beta-(ortho-methoxyphenyl)-isopropyl-methylamine bydrochloride-a bronchodilator and antispasmodic.

*Trademark, Reg. U. S. Pat. Of.

a product of Upjohn

Research

for medicine . . . produced with care . . . designed for health

THE COLORADO STATE MEDICAL SOCIETY

NEXT ANNUAL SESSION: STANLEY HOTEL, ESTES PARK, SEPTEMBER 9, 10, 11, 12, 1952.

OFFICERS

Terms of Officers and Committees expire at the Annual Session in the year indicated. Where no year is indicated, the term is for one year only and expires at the 1952 Annual Session.

President: Harry C. Bryan, Colorado Springs. President-Elect: William A. Liggett, Denver. Vice President: Claude D. Bonham, Boulder.

Constitutional Secretary (three years): Irvin E. Hendryson, Denver, 1954. Treasurer (three years): George C. Shivers, Colorado Springs, 1953.

Additional Trustes (three years): Cyrus W. Anderson, Denver, 1952; E. Munro, Grand Junction, 1952; M. L. Phelps, Denver, 1953; Robert T. Porter, Greeley, 1954.

(The above nine officers compose the Board of Trustees of which Dr.

Board of Councilors (three years): District No. 1: Paul B. Hildebrand, Brush, 1954; No. 2: Kila A. Mead. Greeley, 1954; No. 3: Osgoode S. Philpott, Denver, 1954; No. 4: Ward C. Fenton, Rocky Ford, 1953; No. 5: Jesse W. White, Pueblo, 1953; No. 6: Herman W. Roth, Monte Vista. 1953 (Vice Chairman 1951-1952); No. 7: Leo W. Lleyd, Durango, 1952 (Chairman 1951-1952); No. 8: Harvef M. Tuper, Grand Junction, 1952; No. 9: Marvel L. Crawford, Steamboat Springs, 1952.

Beard of Supervisors (two years): Sidney M. Reckler, Denver, Secretary, 1952; John L. McDonald, Colorado Springs, 1952; Franklin J. McDonald, Leadville, 1952; C. Bex Fuller, Salids, 1953; Lawrence L. Hick, Delta, 1952; Jackson L. Straub, Jr., Flagler, 1952; Lawrence D. Buchanan, Wray, 1953; Jackson L. Sadler, Fort Collins, 1953; Guy C. Carr, Grand Junction, Vice Chairman, 1953; David W. McCarty, Longmont, 1953; V. V. Anderson, Del Norte, 1953; George M. Myers, Pueblo, Chairman, 1953.

Delegates to American Medical Association (two years): William H. Halley, Denver, 1952; (Alternate, Kenneth C. Sawyer, Denver, 1952); Goorge A. Unfug, Pueblo, 1953; (Alternate: Herman C. Graves, Grand Junction, 1953).

Foundation Advocate: Walter W. King, Denver.

House of Delegates: Speaker, Lester L. Ward, Pueblo; Vice Speaker, Kenneth H. Beebe, Sterling.

Exestive Office Staff: Mr. Harvey T. Sethman, Executive Secretary; Miss Helen Kearney, Assistant Executive Secretary; Mr. Evan A. Edwards, Public Relations Director and Field Secretary, 835 Republic Building, Denver 2, Colorado. Telephone AComa 0547,

General Counsel: Mr. J. Peter Nordlund, Attorney-at-Law, Denver.

STANDING COMMITTEES

Arrangaments: Wm. M. Covode, Denver, Chairman; Joseph A. McMeel, Denver; Robert M. Du Roy, Denver; H. P. Thode, Biair Adams, Fort Collins; J. O. Mall, Extes Park; Mr. Harold L. Swanson, Denver.

Credentials: Irvin E. Hendryson, Denver, Chairman; others to be ap-

Health Education (two years): R. A. L. Swanson, Greeley, 1952; Charley J. Smyth, Denver, 1952; W. C. Service, Colorado Springs, 1952; Lewis Barbato, Denver, 1952; W. Lloyd Wright, Golden, 1952; Miss Norma Johannis, Denver, 1952; Doris Benes, Haxtun, 1952; Donald P. Monty, Denver, 1953; Ted W. Miller, Pueblo, 1953; J. D. Bartholomew, Boulder, Chairman, 1953; Gordon Neilgh, Jr., Boulder, E. C. Likes, Lamar; E. Miner Morril, Fort Collins; Paul B. Stidham, Grand Junction.

Library and Medical Literature: Nolie Mumey, Denver, Chairman; Richard Mellen, Colorado Springz; Joel R. Husted, Boulder; W. W. King, Denver; sonard Freeman, Denver.

Medical Education and Mospitals: Cyrus W. Anderson, Denver, Chairman; Marvin Johnson, Denver; Robert S. Liggett, Denver; G. R. Wright, Longmont; Roy F. Dent, Jr., Colorado Springs; Charley J. Smyth, Denver; Robert C. Lewis, Ph.D., Denver; Chaz W. Huff, Jr., Denver; Samuel B.

Medical Service Pinas: Harry C. Hughes, Denver, Chairman; Henry Buchtal, Denver; Charles Gaylord, Longmont; Fredrick H. Good, Denver; Ch. R. Ditsmier, Longmont; V. A. Gould, Mesker; Nathan Beebe, Fort Cullins; Lester L. Ward, Pueblo; Paul G. duSols, Culorado Springs; James R. Blatz, Denver; Alson F. Flerce, Colorado Springs;

Medicelegal (two years): Budolph W. Arndt, Denver, Chairman, 1952; William W. Haggart, Denver, 1952; Edward J. Meister, Denver, 1952; C. S. Bluemel, Denver, 1953; H. I. Barnard, Denver, 1953; E. L. Harvey, Den-

Metrology: C. F. Kemper, Denver, Chairman; Roger S. Whitney, Colorado Springs; C. W. Maynard, Pusblo.

Pablic Policy: Frank B. McGione, Denver, Chairman; Gatewood C. Milli-an, Englewood, Vice Chairman; Wm. B. Candon, Denver; Ervin A. Hinds, server; Karl Arndt, Denver; James Delkous, Denver: V. L. Bolton, Colorado grings; L. D. Buchanan, Wray; Lanning Likes, Lama; Thomas K. Mahan,

Grand Junction; George C. Christie, Canon City; Francis Adams, Pueble D. W. McCarty, Longmont; Harry C. Bryan, Colorado Springs, President Wm. A. Liggett, Denver; Irvin E. Hendryson, Denver, Constitutional Scere

Sub-Committee on Hespital and Prefessional Relations: Ervin A. Hinds, Denver; V. L. Bolton, Colorado Springs; Thomas E. Bost, Denver; Lawrence Campbell, H. J. Dedge, Martin Alexander, John G. Hemming, Pr., George R. J. McDonald, Denver, Chairman; George F. Wigast, Denver; Robert Stere, Denver; Thomas Kennedy, Denver; John Weaver, Jr., Denver.

Sub-Committee on Publicity: Cyrus W. Anderson, Irvin E. Hendryson B. Condon, Ervin A. Hinds, Karl Arndt, Bradford Murphey, all of i

Sub-Committee on Legislation: B. T. Daniels, Denver, Chairman; Karl Aradt, Denver; others to be appointed.

Sab-Committee on Nurses' Education: Walter E. Vest, Jr., Denver, Chairman; John R. Evana, Denver; Carl S. Gydesen, Colorado Springs; Fred D. Kuykendall, Eaton; Miss Mary Walker, Denver.

Sab-Committee on Weekly Healh Column: Howard Bramley, Chairman; Frank Campbell, H. J. Dodge, Martin Alexander, John G. Hemming, Jr., George Curfman, Jr., Charles G. Gabelman, Marianna Gardner, all of Denver.

Subermittee en Farm Magazine Series: Raymond C. Seannell, Denver, Chairman; Paul R. Hildebrand, Brush; William A. Liggett, Denver; Claude D. Bonham, Boulder; David W. McCarty, Longmont; Robert W. Gordon, Denver: Charles A. Rymer, Denver

Scientific Werk: E. Paul Sheridan, Denrer, Chalrman; John C. McAfee, Denrer; Gilbert Balkin, Denrer; E. F. Gerer, Colorado Springs; Felice Garcia, Denver; Kenneth C. Sawyer, Denver; Joseph Lyday, Denrer; J. Cobert Spencer, Mall. Estes Park; Frederick H. Brandenburg, Denver; J. Robert Spencer, Denver; George Curfman, Denver,

PUBLIC HEALTH COMMITTEES

General Committee on Public Health: Consists of the chairmen of the following ten public health sub-committees; presided over by Harold D. Palmer, Denver, as General Chairman.

Caseer Centrel: Harold Palmer, Denver, Chairman; C. B. Kingry, Denver; N. Paul Isbell, Denver; John B. Grow, Denver; R. B. Lanier, Littleton; W. C. Herold, Colorado Springs; C. L. Davis, D.V.M., Denver; J. T. P. Barwick, Pueblo; James A. Philpott, Jr., Denver; Joseph Patternon, Denver; David Akers, Denver; Carl McLauthlin, Jr., Denver; Sidney Reckler, Denver; Mr. Hugh Terry, Denver; Sion W. Holley, Loveland.

Chrenie Diseases: Ward Darley, Denver, Chairman; George A. Unfug, Pueblo; Edward Delehanty, Jr., Denver; Roland A. Raso, Grand Junction; H. B. Haymond, Greeley; Robert Smith, Colorado Springa; Karl J. Waggener, Pueblo; Robert Gordon, Denver.

Industrial Health: James Cullyford, Denver, Chairman; R. H. Ackerly, Pueblo; Robert Bell, Denver; A. R. Woodburne, Denver; Mr. E. W. Jacoe, Denver; Richard C. Vanderhoof, Colorado Springs; James Donnelly, Trinidad, Mr. Ray McBrian, Denver; J. J. Parker, Grand Junction.

Maternal and Child Health: John H. Ameme, Denver, Chairman; E. Stewart Taylor, Denver; Richard K. Kerr, Colorado Springs; Jackson L. Sadder, Fort Collins; Craig Johnson, Denver; L. W. Roessing, Denver; Paul D. Bruns, Denver; John A. Lichty, Denver.

Mental Mysiene: P. H. Zimmerman, Pueblo, Chairman; Bradford Murphey, Denver; E. W. Busse, Denver; Spencer Sayles, Boulder; Warren H. Walker, Denver; Franklin G. Ebaugh, Denver; C. S. Bluemel, Denver; E. James Brady, Colorado Springs; Lewis Barbado, Denver; Clyde Stanfield, Deuver.

Rehabilitation and Crippiod Children: E. L. Binkley, Denver, Chairman; John G. Griffin, Denver; William A. Dorsey, Denver; S. E. Blandford, Jr., Denver; James A. Johnson, Colorado Springs; John C. Long, Denver; Charles G. Freed, Denver; Harold Dinken, Denver; John Bricker, Denver; H. C. Fisher, Denver; M. M. Ginsburg, Denver; Foster Matchett, Denver; Mr. Walter Loague, Denver; Mr. Dorsey Richardson, Denver.

Rural Health and Health Councils: Monroe Tyler, Denver, Chairman; Fred Humphrey, Fort Collins; Robert M. Lee, Fort Collins; Valentin Wohlauer, Akron; H. A. Sauberli, Denver; Kenneth E. Prescott, Grand Junction; John C. Straub, Jr., Flagler; Harlan E. McClure, Lanar; Clement F. Knobbe, Monte Vista; Mr. Low Toyne, Denver; Mr. Marvin Russell, Denver; Mrn. Tee Sims, Denver; Mrs. John Knifton, Sterling; Clara Anderson. Denver.

Sanitation: Bernard T, Daniels, Denrer, Chairman; H. J. Dodge, Denver; Alexis Lubchenco, Denver; Stephen L. Kallay, Denver; Edward N. Chapman, Colorado Springs; W. B. Crouch, Colorado Springs; W. B. Crouch, Colorado Springs; Mr. William Gahr, Denver; Mr. Bobert Cameron, Denver; Mrs. J. W. Penfold, Denver; Man B. Kennon, Denver; Mr. Jean Breitenstein, Denver; Robert Barnard, Aspen; Carl W. Swarts, Fueblo

Teberesiesis Ceetrel: John Zarit, Denver, Chairman; W. J. Hinnelman, Greeley; A. M. Mullett, Colorado Springs; Leroy Eirick, Denver; H. M. Van Der Schow, Wheatridge; Mrs. Ira Waterman, Colorado Springs; Mr. Jack Foster, Denver; Paul B. Marasco, Grand Junction; L. W. Holden, Boulder; Joseph Cannon, Denver; Robert B. Liggett, Denver.

Veneral Disease Control: Sam W. Downing, Denver, Chairman; J. B. McDowell, Denver; Harley Rupert, Greeley; Frederick Tice, Pueble; Joseph Sherman, Denver; Daniel G. Monaghan, Denver.

THE COLORADO STATE MEDICAL SOCIETY

SPECIAL COMMITTEES

Advisory Committee to Woman's Auxiliary: Wiley Jones, Denver, Chairman; McKinnie L. Phelps, Denver,

Advisory to U.M.W. Welfare Fund: W. W. Haggart, Denver, Chairman, 1953; Robert Bell, Denver, 1953; F. H. Hartzhorn, Denver, 1953; J. M. Lamme, Sr., Walsenburg, 1952; Ligon Price, Hayden, 1952; D. W. McCarty, Longmont, 1952; E. E. Ley, Poeblo, 1954; Mason M. Light, Gunnison, 1954; John S. Bouslog, Denver, 1954.

Committee on A.M.A. Educational Campaign: (to be appointed).

Delegate to Colorado interprefessional Council (five years): L. R. Safarik, enver, 1954; J. R. Evans, Denver, 1954, Alternate.

Medical Disaster Commission: Roy L. Cleere, Denver, Chairman; Roger N. Chisholm, Denver; Foster Matchett, Denver; O. S. Philpott, Denver; Harry C. Hughes, Denver; Robert Woodruff, Denver; Karl F. Sunderland, Denver;

Henry Swan, Denver; R. E. Glehm, Denver; Mordant E. Peck, Denver; M. P. Vanden Bosch, Denver; T. P. Sears, Fort Logan; M. E. Johnson, Denver; B. L. Goldman, Denver; Mark S. Donoran, Denver; McGordick J. McDonald, Denver; M. B. Pedigo, Denver; W. S. Curtis, Denver; K. D. A. Allen, Denver; K. E. Gloss, Colorado Springs.

Military Affairs Committee: Robert S. Liggett, Denver, Chairman; Calvin N. Caldwell, Pueblo; Ward C. Fenton, Rockly Ford; Leo W. Lloyd, Durango; Frank I, Nicha, Colorado Spring; Claude D. Bonham, Boulder; Harvey M. Tupper, Grand Junction; George E. Buck, Denver; John P. Foster, Denver.

Representative to Rocky Mountain Radio Council: Irvin E. Hendryson.

Representatives to Adult Education Council: Cyrus W. Anderson, Denver; William E. Hay, Denver.

Rocky Mountain Medical Conference: G. P. Lingenfelter, Denver, Chairman, 1952; D. W. Macomber, Denver, 1954; L. Clark Hepp, Denver, 1953; Terry J. Gromer, 1955; William Covode, Denver, 1956.

Two Special DAIRY FOODS which physicians can prescribe confidently

For persons OVER-WEIGHT or on a LOW-FAT Diet

HIGH in Vitamins LOW in Calories

Butterfat removed — Vitamins added (4,000 units Vitamin A, 400 units Vitamin D). 88 calories per quart.

"At Your Store or



For persons UNDER-WEIGHT needing Extra Nutrition

GOLDEN GUERNSEY

Contains 4.4 butterfat-with proportionately higher content of milk's important nutrients. Cream-top or homogenized.

CARLSON-FRINK at Your Door"

Denner's Quality Dairy



Oneonta Gorge, Oregon

Special Morning Milk is an evaporated milk especially developed for infant feeding. It is fortified (from the natural source) with 400 U.S.P. units vitamin D and 2000 U.S.P. units vitamin A per reconstituted quart.

> rom the natural source

Special Morning Milk Morni





MONTANA MEDICAL ASSOCIATION

NEXT ANNUAL SESSION: MISSOULA, SEPTEMBER 18, 19, 20, 21, 1952

OFFICERS, 1951-1952

Terms of Officers and Committees expire at the Annual Session in the year indicated. Where no year is indicated, the term is for one year only and expires at 1952 Annual Session.

President: Frank L. McPhail Great Falls President-Elect: James M. Flinn, Helena, Vice President: B. C. Farrand, Jordan,

Secretary-Treasurer: E. H. Lindstrom, Helena.

Asst. Secretary-Treasurer: W. J. Roberts, Great Falls.

Executive Secretary: Mr. L. R. Hegland, 240 Stapleton Bldg., Billings, Delegate to American Medical Association: Raymond F. Peterson, Butte; ternate, Thomas L. Hawkins, Helena.

STANDING COMMITTEES

Executive Committee: F. L. McPhall, Chairman, Great Falls; B. C. Farrand, Jordan; James M. Flinn, Helena; Clyde H. Fredrickson, Missoula; Thomas L. Hawkins, Helena; Everett H. Lindstrom, Helena; Wyman J. Roberts, Great Falls.

Economic Committee: D. Ernest Hodges, Chairman, Billings; R. L. Casebeer, Butle; William F. Cashmore, Helena; William E. S. Harris, Livingston; Robert J. Holzberger, Great Falls; Duncan S. MacKenzie, Jr., Harre; Sidney C. Prati, Miles City; James A. Mueller, Lewistown.

Legislative Committee: I. J. Bridenskine, Chairman, Missoula; Sidney A. noney, Helena; Otto G. Klein, Helena; James J. McCabe, Helena; Richard. Monahan, Butte; Robert M. Morgan, Helena; E. S. Murphy, Missoula; uart D. Whetstone, Cut Bank,

Necrology and Histery of Modicine Committee: Leonard W. Brewer, Chairman, Missoula; Melville G. Danakin, Glendive; Albert A. Dodge, Kalispell; Edward M. Gans, Harlowton; W. G. Richards, Billings; John Paul Ritchey, Missoula; J. I Wernham, Billings; S. V. Wilking, Butte.

Public Relations Committee: Albert W. Axley, Chairman, Havre; Paul L. Enchoe, Bozeman; F. S. Marks, Billings; Arthur K. Northrop, Great Falls; Stuart A. Olson, Glendive; R. F. Peterson, Butte; C. R. Svere, Missoula; Park W. Willis, Jr., Hamilton.

Legal Affairs and Malpraetice Committee: Louis W. Allard, Chairman, Billings; J. H. Bridenbaugh, Billings; H. W. Gregg, Butte; P. E. Logan, Great Falls; T. R. Vye, Billings.

Program Committee: Mary E. Martin, Chairman, Billings; Charles B. Craft, Bozeman; John A. Layne, Great Falls; Stephen N. Preston, Missoula; T. W. Saam, Butte; Everett H. Lindstrom, Helens, Ex-Officio.

Interprofessional Selations Committee: M. A. Shillington, Chairman, Glendive; Louis W. Allard, Billings; J. K. Colman, Butte; Theodore W. Cooney, Helena; Carl W. Hammer, Bozeman; George W. Sexton, Great Falls.

Nominating Committee: G. W. Setzer, Chairman, Malta; Neil M. Leitch, Kalispell; T. R. Vye, Billings; Edmund A. Welden, Lewistown; Malcolm D. Winter. Miles City.

Auditing Committee: George G. Sale, Chairman, Missoula; J. M. Brooke, onan; George M. Donich, Anaconda; Robert D. Knapp, Wolf Point; G. pron Wright, Kalispell.

Caneer Committee: Raymond E. Benson, Chairman, Biltings; Walter B. Cux, Missoula; Deane C. Epler, Bozeman; H. W. Gregg, Butte; E. Hildebrand, Great Falls; K. E. Markuson, Helena, Ex-Officio; Philip D. Pal-

Maternal and Child Welfare Committee: Earl L. Hall, Chairman, Great

Falis.

Sabcommittee on Obstetries: G. A. Carmichael, Chairman, Missoula; Joe E. Brann, Kalispell; Harry B. Campbell, Missoula; Maude M. Gerdos, Billings; G. W. Pemberton, Butte.

Subcommittee on Pediatries: Grville M. Moore, Chairman, Helena; George H. Barmeyer, Missoula; Roger W. Clapp, Butte; Frank J. Friden, Great Falis; D. L. Gillespie, Butte; R. Wynne Morris, Helena; George W. Nelson, Billings; Paul R. Ensign, Helena, Ex-Offico.

Tubercalosis Committee: H. V. Gibson, Chairman, Great Falls; L. M. Arthur, Great Falls; J. K. Colman. Butte; Charles B. Craft, Boocman; Morris Alan Gold, Butte; J. M. Nelson, Missoula; Stephen N. Freston, Missoula; R. E. Smailer, Billings; Frank I. Terrill, Galen; William F. Kimmell, Helena, Ex-Office.

Fracture and Orthopedic Committoe: Walter H. Hagen, Chairman, Billings; L. Clayton Allard, Billings; J. K. Colman, Butte; C. F. Honeycutt, Missoula; S. L. Odgers, Missoula; John A. Whittinghill, Billings; John C. Wolgamot, Great Falls; Paul R. Ensign, Helena, Ex-Officio.

Rural Health Committee: B. C. Farrand, Chairman, Jordan; David Gregory, Glasgow; James M. Isbister, Plains; Burlon K. Kilbourne, Hardin; Robert H. Leeds, Chinook; Ronald E. Louee, Eunis; Walter G. Tanglin, Polson; Amos R. Little, Helcna; George E. Trobough, Anacenda; Lester S. McLean, Helena, Ex-Officio.

Industrial Weifare Committee: R. B. Richardson, Chairman, Great Falls; W. Gregg, Butte: John J. Malee, Anaconda; W. F. Morrison, Missoula; darey C. Fratt, Miles City; George G. Sale, Missoula; James G. Sawyer, ttte; John W. Schubert, Lewistown; F. K. Waniata, Great Falls; K. E.

Rheemstle Fever and Heart Committee: F. R. Schemm, Chairman, Great Falls; Raymond L. Eck, Lewistown; D. L. Gillespie, Butle; John Gilbox, Great Falls; Morris Alan Gold, Butle; Elizabeth Grünn, Billings; C. S. Meeker, Butle; Orville M. Moore, Helena; Thomas F. Walker, Jr., Great Falls; Richard D. Weber, Missouli; G. D. Carlyle Thompson, Helena, Ex-

Rocky Mountain Medical Conference Committee: Chairman, '53; H. M. Blegen, Missoula, '55; H. T. Caraway, Billings, '64; Charles B. Craft, Bozeman, '56; F. K. Waniata, Great Falls, '52; F. L. McPhail, Great Falls, Ex-Officio; Everett H. Lindstrom, Helena, Ex-Officio

Mediation Committee: F. S. Marko, Chairman, Billings, '54; Eamer P. Higgins, Kalispell, '54; Chester W. Lawson, Havre, '52; Charles F. Little, Great Falls, '53; William E. Long, Anaconda, '53; James J. McCabe, Helena, '54; W. F. Morrison, Missoula, '52; Stuart A. Olsen, Glendive, '53; James G. Sawyer, Butte, '52.

James G. Sawyer, Sutze, '52.

Pablic Health Committee: James M. Flinn, Chairman, Helena; Raymond E. Benson, Billings; Deane C. Epler, Bozeman; B. C. Farrand, Jordan; H. V. Gibson, Great Falls; Walter H. Hagen, Billings; Earl L. Hall. Great Falls; E. Hildebrand, Great Falls; Amos R. Little, Helena; R. B. Richardson, Great Falls; F. R. Schemm, Great Falls; M. A. Shillington, Glendüre; Walter G. Tanglin, Polson; George E. Trobough, Anaconda; Winfield S. Wilder, Great Falls.

SPECIAL COMMITTEES

Emergency Medical Service Committee: Amos R. Little, Chairman, Helena; David J. Almas, Harre; Leonard M. Benjamin, Deer Lodge; Leonard W. Benyamin, Deer Lodge; Leonard W. Brewer, Missoula; Harrison D. Huggdina, Kalispelj, Leland G. Russell, Billings; H. J. Sannan, Butte; Philip A. Smith, Glasgow; Albert L. Vadheim, Boseman; Thomas F. Walker, Jr., Great Falls; G. D. Carlyle Thompson, Helena, Ex-Officio.

Hospital Relations Committee: E. Hildebrand, Chairman, Great Falls; beert B. Beans, Great Falls; Waiter B. Cox, Missoula; E. W. Gibbe, Illings: Robert S. Leighton, Great Falls; Mary E. Martin, Billings; W. W. cLaughlin, Great Falls; R. F. Peterson, Butte; F. M. Petkerich, Great Falls: Grant P. Raitt, Billings.

Mental Hypiene Committee: Winfield S. Wilder, Chairman, Great Falls; James J. Bulger, Great Falls; Boger W. Clapp, Butte; G. V. Holmes, Missoula; J. E. Kress, Missoula; Martin A. Buona, Billings; M. A. Shillington, Glendire.


Physicians-Schools Conference: Ray O. Bjork, Chairman, Helena; George M. Donich, Anaconda; Earl L. Hall, Great Falls; Eaner P. Higgins, Kalispell; Stuart A. Olson, Glendive; C. R. Svore, Missoula.

Revision of By-Laws Committee: Thomas L. Hawkins, Chairman, Helena; Paul J. Gans, Levistown; Eaner P. Higgins, Kalispell; Wyman J. Roberts, Great Falis; M. A. Shillington, Glendive.



Cambridge Dairy Grade "A" Milk Is Produced and Processed at 690 S. Colo. Blvd. We do not handle Shipped-in Milk produced Where? How and by Whom? Doctors know the difference Now Homogenized Vitamin D Milk is available for baby feeding and family use, We Invite Your Inspection and Appreciate Your Recommendation.

М١



Psychoneurotics of Genius

Hawthorne, distinguished American novelist, is said to have been afflicted with a psychoneurosis from early childhood. His quiet life, wholly detached from the major activities of the times, was largely given over to bro

The majority of psychoneurotics have no serious mental illness, but display merely an emotional imbalance which often can be greatly improved by appropriate psychotherapeutic and sedative management. In the treatment of psychoneurosis, particularly agitated, depressed and anxiety states, Mebaral is especially useful when tranquillity with minimal hypnotic action is desired. Sedative dose: Adults, from 32 mg. to 0.1 Gm. (½ to 1½ grains) three or four times daily. Children, from 16 to 32 mg. (½ to ½ grain) three or four times daily, Supplied in tablets of 32 mg., 0.1 Gm. and 0.2 Gm.

MEBARAL® Brand of Mephobarbital

Tasteless SEDATIVE AND ANTIEPILEPTIC
Little or No Drowsiness

WINTHROP-STEARNS INC. . NEW YORK 18, N. Y. . WINDSOR, ONT.

Naharal, trademark rea, U.S. & Conada

NEW MEXICO MEDICAL SOCIETY

NEXT ANNUAL SESSION: CARLSBAD, MAY 8, 9, 10, 1952

OFFICERS—1951-52

President: Leland S. Evans, Las Cruces President-Elect: Coy S. Stone, Hobbs.

Vice President: Albert S. Lathrop, Santa Fe.

Secretary-Treasurer: L. G. Rice, Jr., 611 East Central, Albuquerque Executive Secretary: Mr. Ralph Marshall, 323 First National Bank,

Councilors (3 years): W. D. Dabbs, Clovis; W. E. Badger, Hobbs. (2 years): A. S. Lathrop, Santa Fe; Carl H. Gellenthien, Valmora. (1 year): Carl Mulky, Albuquerque; J. C. Sedgwick, Las Croces.

New Mexico Physicians' Service: President, John F. Conway, Clovis; Vice President, Victor K. Adams, Raton; Secretary-Treasurer, L. G. Rice, Jr., Albuquerque; Executive Director, Mr. Leuis J. LaGrave, Box 1082, Albu-

Board of Trustees: L. S. Erans, Las Cruces; A. H. Follingstad, Albuquerque; Carl H. Gellenthien, Valmora; Albert S. Lathrop, Santa Fe; H. A. Miller, Clovis; George S. Morrison, Clovis; Ashley C. Shuler, Carlsbad; W. A. Stark, Las Vegas.

COMMITTEES-1951-52

Beard of Supervisors (Two Years): H. M. Mortimer, Las Vegas; Earl L. Malone, 113 North Kentucky, Rousell, Sceretary; L. J. Whitaker, Dening; Prank W. Patker, Galluo, (One Year): C. Pardos Bunch, Artesia, Chairman; H. L. January, Albuquerque; John F. Conway, Clovis; V. E. Berchtold. Santa Fe, Vice Chairman.

Basic Science Committee: Marcus J. Smith, Santa Pe, Chairman; Brian Moynahan, Albuquerque; W. D. Anthony, Gallup.

Cancer Committee: Charles Moreau Thompson, Albuquerque, Chairman;
J. W. Grossman, Albuquerque; Aaron E. Margulis, Santa Fe; Murray M. Priedman, Santa Fe; M. Waggoner, Roowell; Loren Blaney, Los Alamos, Diabetic Committee: John H. Dettweller, Albuquerque, Chairman; Alfred, J. Jemson, Hobbs; Bergsre A. Kenney, Santa Fe; J. E. Merritt, Las Cruces, Lefaney, Company, Company

infancy and Maternal Care Committee: Alfred C. Service, Roswell hairman; Lee M. Miles, Albuquerque; Oscar Syme, Albuquerque; S. M. onzales, Santa Fe; Charles E. Galt, Carlsbad; Marion Hotopp, Santa Fe.

industrial Health Committee: Lewis M. Overton, Albuquerque, Chairman; U. S. Marshall, Roswell; Edgar A. Rygh, Santa Rita; J. H. Burress, Raton; J. W. Hillsman, Carlsbad; N. D. Fraxin, Silver City.

indigent-Medical Care Committee: A. C. Rood, Albuquerque, Chairman; Samuel R. Ziegler, Espanola; J. J. Johnson, Les Vegas.

Samuer R. Ziegier, Espanola; J. J. Jonnson, Les Vegas.

Lesislative and Pablic Polley Cosmittee: A. S. Lathrop, Santa Fe,
Chairman; J. W. Hannett, Albuquerque; Milton Floersheim, Raton; V.
Scott Johnson, Clovis; L. L. Daviet, Las Cruces; A. T. Gordon, Tucuewart;
Martin S. Withers, Loe Alamos; Clay A. Gwinn, Carisbad; Junius A. Evans,
Las Vegas; Charles F. Kettel, Gallup; W. L. Minear, Truth or Consequences;
R. E. Watts, Silver City; Ashley Pond, Taon; Cop S. Stone, Hobby; W. J.
Hossley, Deming; I. J. Marshall, Roswell; Wesley O. Conner, Jr., Albu-

National Emergency Nedical Service Committee: Andrew J. McQueeney, Albuquerque, Chairman; William R. Oakes, Los Alamos; Richard A. Angle, Santa Fe; W. A. Stark, Los Vegas; H. O. Lehman, Portales; Samuel M. Ramer, Sliver City.

Public Relations Committee: E. C. Derbyshire, Santa Fe, Chairman; Earl L. Malone, Roswell; Hilton W. Gillett, Lovington; George W. Prothro, Clovis; W. D. Bedgwick, Las Cruces.

Raral Health Committee: Benjamin Barzune, Eunice, Chairman; Stuart W. Adler, Albuquerque; J. P. Turner, Carrizozo; Wendell H. Peacock, Farmington; C. E. Molholm, Grants; Eugene P. Shams, Alamogordo.

Tukercaiesis Committee: Carl H. Gellenthien, Valmora, Chairman; W. H. Thearle, Albuquerque; Carl Mulky, Albuquerque; H. C. Jernigan, Albuquerque; H. S. A. Alexander, Santa Fe.

querque; H. S. A. Alexander, Santa Fe.

Vanerasi Dissase Control Committee: H. J. Beck, Albuquerque, Chairman;
T. E. Kircher, Jr., Albuquerque; I. L. Feavy, Santa Fe; David T. Wier,
Belen; J. H. Donnelly, Portales.

Woman's Auxiliary Committee: Philip L. Travers, Santa Fe, Chairman;
Louis A. McRae, Albuquerque; W. N. Worthington, Rowell.

Rocky Meantain Medical Conference Committee: Carl H. Gellenthien,
Valmora, Chairman; Victor E. Adams, Raton; J. W. Beattie, Las Vegas;
Eric P. Hausner, Santa Fe; A. H. Follingstad, Albuquerque.

Eye and Ear Consulting Committee to State Dept. of Pablic Health:
Howard B. Peck, Albuquerque; George S. Richardson, Albuquerque; R. R.
Boice, Rowell; James L. McCroory, Santa Fe; A. W. Egenhofer, Santa Fe
Advisory Committee on Insurance Compensation: R. W. McIntyre, Albuquerque, Chairman; Gerald A. Siusser, Silver City; Peter J. Starr, Artesia.



Our dairy farm is the largest producer of Grade "A" milk in the Rocky Mountain Empire.

EAst 7707

CITY PARK FARM DAIRY Cherry Creek Dr.



FAIRFAX SANITARIUM


Kirkland, Wash.

Situated one mile north of Juanita TREATING NERVOUS AND MENTAL DISEASES

Beautiful and restful surroundings affording recreational facilities. Cottage plan for segregation of patients. Insulin and Electro-shock Therapy when indicated.

Attending Physicians Attending Physicians
FREDERICK LEMERE, M.D.
NATHAN K. RICKLES, M.D.
JAMES H. LASATER, M.D.
MORTON E. BASSAN, M.D.
JACK J. KLEIN, M.D. Manager: A. G. HUGHES

Route 2, Box 365, Kirkland Phone: Kirkland 2391



PETROGALAR provides a moderate intake of mineral oil in the form of a water-miscible suspension.

This oil-in-water combination permeates the fecal residue to produce:

- ► Gentle lubricant action, without "leakage"
- Soft, nonirritating, easily passed stools
- Comfortable bowel movement

PETROGALAR may be taken alone or in milk, water or fruit juices—with which it is readily miscible.

PETROGALAR®

Aqueous Suspension of Mineral Oil, Wyeth

Wyeth Incorporated, Philadelphia 2, Pa.



THE UTAH STATE MEDICAL ASSOCIATION

NEXT ANNUAL SESSION, SALT LAKE CITY, SEPTEMBER 4, 5, 6, 1952

OFFICERS, 1951-52

President: L. W. Oaks, Provo.

President-Elect: Kenneth B. Castleton, Salt Lake City.

Past President: V. P. White, Salt Lake City.

Nonerary President: Jos. R. Morrell, Ogden.

First Vice President: R. P. Middleton, Salt Lake City. Second Vice President: C. C. Bandall, Logan.

Third Vice President: P. R. King, Price. Secretary: T. C. Weggeland, Salt Lake City.

Executive Secretary: Mr. W. H. Tibbals, Salt Lake City.

Treasurer: L. J. Paul. Salt Lake City.

Treasurer: L. J. Paul, Salt Lake City.

Councilor, First District: R. O. Porter, Logan.

Councilor, Second District: Vincent L. Rees, Salt Lake City.

Councilor, Third District: J. Russell Smith, Provo.

Delegate to A.M.A., 1952 and 1953: George M. Fister, Ogden. Alternate Delegate to A.M.A., 1952 and 1953: J. J. Weight, Provo.

Editor of the Utah Section of the Rocky Mountain Medical Journal: R. P. Middleton, Salt Lake City.

Beard of Sepervisors: 1952, Paul K. Edmunds, Cedar City; 1953, Earl L. Skidmore, Salt Lake City; 1954, J. C. Hubbard, Price; 1955, J. G. Olson, Ogden; 1956, C. J. Daines, Logan.

STANDING COMMITTEES

Rocky Mountain Medical Conference Continuing Committee: 1952, Noall Z. Tanner, Chairman, Layton; 1953, T. R. Seager, Vernal; 1954, R. P. Middleton, Salt Lake City; 1955, U. R. Bryner, Salt Lake City; 1956, Heber C. Hancock, Ogden.

Scientific Program Committee: T. C. Weggeland, Chairman, Salt Lake City.

Public Policy and Legislative Committee: 1952, Charles Ruggerl, Chairman, Salt Lake City; 1952, J. C. Rubbard, Price; 1952, Wilford G. Blesinger, Springville: 1953, N. F. Hicken, Salt Lake City; 1953, L. V. Broadbent, Cedar City; 1953, George Gasser, Logan; 1954, V. L. Stevenson, Sait Lake City; 1954, Charles R. Cornwall, Salt Lake City; 1954, John Z. Bowers, Sait Lake City; Wendell Thomson, Ogden; Claude L. Shields, Salt Lake City; R. M. Muirhead, Salt Lake City; C. Eliot Snow, Salt Lake City; Roy B. Hammond, Provo; Conrad H. Jenson, Ogden; Ralph Richards, Salt Lake City.

Sub-Committee on Legislation: Vernon L. Stevenson, Chairman, Salt Lake City; George Gasser, Logan; Charles R. Cornwall, Salt Lake City; L. V. Broadbent, Cedar City; John Z. Bowers, Salt Lake City; N. F. Hicken, Salt Lake City.

Medical Defense Committee: 1952, E. L. Hanson, Logan; 1952, Reed Farnsworth, Cedar City; 1952, H. A. Dewey, Richfield; 1953, John B. Cluff, Richfield; 1953, Paul A. Pemberton, Salt Lake City; 1953, Wendell Thomson, Ogden; 1954, R. W. Owens, Chairman, Salt Lake City.

Medical Education and Nospitals Committee: 1952, Ralph Ellis, Ogden; 1952, Philip Price, Salt Lake City; 1952, W. H. Anderson, Ogden; 1953, T. C. Bauerlein, Salt Lake City; 1953, E. R. Crowder, Salt Lake City; 1953, E. R. Crowder, Salt Lake City; 1954, Harry J. Brown, Chairman, Provo; 1954, L. K. Gates, Logan; 1954, K. A. Crockett, Salt Lake City; 1955, R. V. Larzen, Rosserelt; 1965, Mark B. Jensen, Castle Gate; 1955, J. B. Cluff, Richfield; 1954, W. J. Reichman, St. George; John M. Waldo, Salt Lake City.

Sub-Committee on Postgraduate Education: R. V. Larsen, Chairman, Roosevelt; Mark B. Jensen, Castle Gate; J. B. Cluff, Richfield; W. J. Reichman, St. George; John M. Waldo, Salt Lake City.

Medical Economics Committee: 1952, Grant F. Kearns, Ogden; 1952, Preston Hughes, Spanish Fork; 1953, Hugh O. Brown, Chairman, Salt

Lake City; 1953, Silas S. Smith, Salt Lake City; Ralph N. Barlow, Logan.

Public Health Committee: 1952, R. N. Hirst, Ogden; 1952, James Z. Davis, Salt Lake City; 1953, Paul Clayton, Chairman, Salt Lake City; 1953, Alma Nemir, Salt Lake City; 1953, John Bourne, Provo; 1953, Michael E. Murphy, Salt Lake City; 1953, John Bourne, Provo; 1953, Michael E. Murphy, Salt Lake City; 1953, A. A. Jenkins, Salt Lake City; 1953, John Bowen, Provo; 1954, E. M. Kilpatrick, Salt Lake City; 1954, Preston Catter, Salt Lake City; 1954, Prest Octater, Salt Lake City; 1954, Drew M. Peterson, Ogden; J. H. Bupper, Provo; D. O. N. Lindberg, Ogden.

Sub-Committee on Tuberculosis and Cardiovascular Diseases: E. M. Kilpatrick, Chalrman, Salt Lake City; Preston Cutler, Salt Lake City; Fred W. Clauson, Salt Lake City; Drew M. Peterson, Ogden; J. H. Rupper, Provo; D. O. N. Lindberg, Ogden,

Cancer Committee: John H. Carlquist, Chairman, Salt Lake City; Wm. H. Moretz, Salt Lake City; Angus K. Wilson, Salt Lake City; E. D. Zeman, Ogden; Riley G. Clark, Provo.

Fracture Committee: L. N. Ossman, Chairman, Salt Lake City,

Necrology Committee: L. A. Stevenson, Chairman, Salt Lake City.

Industrial Health Committee: F. J. Winget, Chairman, Sait Lake City; B. F. Robison, Sait Lake City; E. Wayne Alfred, Orem; Noal Z. Tanner, Latyson: Chester Powell, Sait Lake City; R. R. Robinson, Sait Lake City; Wendell Thompson, Ogden; George A. Spendiore, Sait Lake City;

Advisory Committee to the Woman's Auxiliary: L. W. Oaks, Chairman, Provo; Kenneth B. Castleton, Salt Lake City; V. P. White, Salt Lake City; T. C. Weggeland, Salt Lake City; L. J. Paul, Salt Lake City; R. O. Porter, Logan; Vincent L. Rees, Salt Lake City; J. Russell Smith, Provo.

Public Relations Committee: Dean Spear, Chairman, Salt Lake City; R. W. Farnsworth, Cedar City; John Z. Bowers, Salt Lake City; N. F. Hicken, Salt Lake City; George Ely, Salt Lake City; T. R. Seager, Vernal.

Mental Health Committee: Roy A. Darke, Chairman, Salt Lake City; L. G. Moench, Salt Lake City; W. D. O'Gorman, Ogden; O. P. Heninger, Provo; C. H. Branch, Salt Lake City; Lyman Horne, Salt Lake City; F. F. Hatch, Salt Lake City; F. F.

Rural Health Committee: R. W. Farnsworth, Cedar City; L. H. Merrill, Hiawatha; Theodore Noehren, Salt Lake City; John R. Martineau, Morgan.

Sub-Committee Postgraduate Education Committee: B. V. Larsen, Chairman. Roosevelt; Mark B. Jensen, Helper; J. B. Cluff, Richfield; W. J. Reichman, St. George.

Procurement and Assignment Committee: C. Eliot Snow, Chairman, Salt Lake City; Frank K. Bartlett, Ogden; John J. Galligan, Salt Lake City; John H. Clark, Salt Lake City; J. Bussell Smith, Provo.

Civilian Defense Committee: L. J. Paul, Chairman, Salt Lake City; Leo W. Benson, Ogden; Biley G. Clark, Provo; S. M. Budge, Logan; Boyd Larsen, Lehi.

Fee Schedule Committee: W. R. Rumel, Chairman, Salt Lake City; F. F. Hatch, Salt Lake City; John H. Clark, Salt Lake City; Leroy Smith, Salt Lake City; Junior Rich, Ogden; L. N. Ossman, Salt Lake City; R. R. Robinson, Salt Lake City; Scott Smith, Salt Lake City; Chester B. Fowell, Salt Lake City; M. L. Crandall, Salt Lake City; Wm. R. Young, Salt Lake City; Wm. J. Morginson, Salt Lake City; Dean A. Moffat, Salt Lake City; Robert W. Ogilvie, Salt Lake City.

Constitution and By-Laws Committee: Louis P. Matthei, Chairman, Ogden; Kenneth A. Crockett, Sait Lake City; Rulon Howe, Ogden; Irving Ershler, Sait Lake City; Byron Daynes, Sait Lake City; Ray T. Woolsey, Sait Lake

Special Committee to Investigate the Nursing School at Logan, Utah: J. C. Haward, Logan; R. M. Muirhead, Salt Lake City; J. R. Miller, Salt Lake

Gerontology Committee: Richard P. Middleton, Chairman, Salt Lake City.

PROFESSIONAL MEN RECOMMEND

Professional PHARMACY, INS.

D. MALCOLM CAREY, Pharmacist

Phone AComa 3711

224 Sixteenth Street

Denver. Colo.

Better Flowers at Reasonable Prices
"Orders Delivered to Any City by
Guaranteed Service"

Special attention given to floral tributes
Also Hospital Flowers

Call KEystone 5106

Park Floral Co. Store

1643 Broadway

Denver, Colo.



and in peritonitis. For hospital use only.

Supplied 10 cc. vial, 250 mg.;
20 cc. vial, 500 mg.

Terramycin is also available as Capsules, Elixir, Oral Drops, Ophthalmic Ointment, Ophthalmic Solution.

ANTIBIOTIC DIVISION



CHAS. PFIZER & CO., INC., Brooklyn 6, N. Y.

THE WYOMING STATE MEDICAL SOCIETY

NEXT ANNUAL SESSION, LANDER, JUNE 5, 6, 7, 1952

President: Paul R. Holtz, Lander President-Elect: Edward Guilfoyle, Newcastle, Vice President: James Sampson, Sheridan. Secretary: G. W. Koford, Cheyenne. Treasurer: P. M. Schunk, Sheridan.

Executive Secretary: Arthur B. Abbey, Cheyenne. Delegate to A.M.A.: Roscoe H. Reeve, Casper.

Alternate Delegate to A.M.A.: W. Andrew Bunten, Cheyenne.

Rocky Maontain Medical Conference: Earl Whedon, Chairman, Bheridan; George H. Phelps, Cheyenne; H. L. Harrey, Camper; L. W. Storey, Laramie, Syphilis Committee: L. H. Wilmoth, Chairman, Lander; F. H. Haigler, Casper; Benjamin Gittlix, Thermopolis

Cancer Committee: John Gramlich, Chairman, Cheyenne; Benjamin Gitlitz, Thermopolis; Thomas B. Croft, Lovell; Karl E. Krueger, Rock Springs; Franklin Yoder, Cheyenne.

Medical Economies Committee: Ernest A. Kahn, Chairman, Cheyenne; George M. Knapp, Casper; Carleton D. Anton, Sheridan.

Fracture Committee and Industrial Health: Gordon Whiston, Chairman, Camper; DeWitt Dominick, Cody; E. C. Pelton, Laramie; Paul Preston, Cheyenne.

Advisory Committee to Selective Service on Procurement and Assignment of Physicians: Sam S. Zuckerman, Chairman, Cheyenne; Roscoe H. Roeve, Casper; E. W. DeKay, Laramie.

Elected: Medical Defense Committee: DeWitt Dominick, Chairman, Cody;
A. Bunten, Cheyenne; Karl E. Krueger, Rock Springs.

Councillors: Karl E. Krueger, Chairman, Rock Springs; Earl Whedon, Sheridan; George Baker, Casper; DeWitt Dominick, Cody: George H. Phelps, Cheyenne; Paul R. Holts, President, Lander; Glean W. Koford, Secretary.

Advisory to Woman's Auxiliary: J. Cedric Jones, Chairman, Cody; John R. Bunch, Laramie: W. A. Bunten, Cheyenne.

Veterans Affairs and Military Service Committee: Bernard Sullivan, Chairman, Laramie; R. C. Stratton, Green River; James Sampson, Sheridan; A. J. Allegretti, Cheyenne; E. J. Guilfoyle, Newcastle.

Bise Cross Hespital Committee: Russell Williams, Chairman, 1954, Cheyenne; E. W. DeKsy, 1955, Laramie; DeWitt Dominick, 1953, Cody; J. W. Sampson, 1953, Sheridan.

Public Policy and Logislation: G. W. Koford, Chairman, Cheyenne; corge H. Phelps, Cheyenne; W. A. Bunten, Cheyenne; E. W. DeKay, aramie; Paul R. Holtz, Lander; R. H. Reere, Casper.

Pollomyelitis Committee: L. Cohen, Chairman, Cheyenne; E. W. Gardner, uglas; Franklin Yoder, Cheyenne; M. C. Henrich, Casper.

State Institutions Advisory: R. H. Kanable, Chairman, Besin; Franklin Yoder, Cheyenne; Joseph F. Whalen, Evanston; L. H. Wilmoth, Lander. Nearlogy Committee: Earl Whedon, Chairman, Sheridan; Franklin Yoder, Cheyenne.

Public Health Department.—Linison Committee: E. C. Ridgway, Chairman, Cody: R. P. Fitzgerald, Casper; J. W. Sampson, Sheridan; R. C. Stratton, Green River.

Rural Health Committes: Andrew Bunten, Chairman, Cheyenne; William Rosene, Wheatland; Samuel H. Worthen, Afton; John B. Krahl, Ter-

Child Health Committee: Paul Emerson, Chairman, Cheyenne; L. J. shen, Cheyenne; M. C. Henrich, Casper,

Council on National Emergency Medical Service: George H. Fhelps, Chairman, Cheyenne; R. H. Reeve, Casper; E. W. DeKay, Laramie; P. M. Schunk, Sheridan; Paul R. Holts, Lander; Albert T. Sudman, Green River; DeWitt Dominick, Cody.

Committee for Professional Review: J. D. Shingle, Chairman, Cheyenne; Paul R. Holts, Lander; J. Cedric Jones, Cody; Gordon C. Whiston, Casper.

COLORADO HOSPITAL ASSOCIATION

OFFICERS

President: Henry H. Hill, Weld County Hospital, Greeley.

President-Elect: H. E. Rice, Porter Sanitarium and Hospital, Denver.

Vice President: Sr. Marie Charles, Glockner-Penrose Hospital, Colorado

Treasurer: M. A. Moritz, Denver General Hospital, Denver.

Executive Secretary: R. A. Pontow, Colorado General Hospital, Denver.

Trastees: Roy R. Prangley, St. Luke's Hospital, Denver (1952); James P. Dixon, M.D., Denver General Hospital, Denver (1953); G. A. W. Currle, M.D., Colorado General Hospital, Denver (1954); Louis Lisseod, National Jewish Hospital, Denver (1952); A. Tergerson, Longmont Hospital, & Clinic, Inc., Longmont (1953); DeMoss Tallaferro, Children's Hospital, Denver (1954).

Delegate to American Hospital Association: Msgr. John R. Mulrey, Catholic Hospitals, Denver.

Alternate: Herbert A. Black, M.D., Parkview Hospital, Pueblo,

COMMITTEES FOR 1952

Auditing: Ed Smith, Chairman, Boulder Colorado Sanitarium and Hospital, Boulder (1952); John Peterson, Larimer County Hospital, Fort (1953); Paul Tadlock, Colorado General Hospital, Denver (1954).

Legislative: Hubert Hughes, Chairman, General Rose Memorial Hospital, Denver; Magr. John R. Muiroy, Catholic Hospitals, Denver; DeMoss Tallaterro, Children's Hospital, Denver; Roy Anderson, Presbyterian Hospital, Denver; F. H. Zimmerman, M.D., Colorado State Hospital, Pueblo.

Membership: Louis Lismood, Chairman, National Jewish Hospital, Denver; A. Tergerson, Longmont Hospital and Clinic, Inc., Longmont; Sister M. Ascella, St. Joseph's Hospital, Denver.

Nominating: Magr. John R. Mulroy, Chairman, Catholic Hospitals, Den-r (1953); A. Tergerson, Longmont Hospital and Clinic, Inc., Longmont

Nursing Education: Roy R. Frangley, Chairman, St. Luke's Hospital, Denver; Slster M. Hugolina, St. Anthony's Hospital, Denver; Marguerite E. Factznick, Denver General Hospital, Denver; Rev. Allen Erb, Mennonito Hospital and Sanitarium, La Junta; Mrs. Henrietta Loughran, University of Colorado School of Nursing, Denver.

Program: H. E. Rice, Chairman, Porter Sanitarium and Hospital, Denver; Charles K. Levine, Beth Israel Hospital, Denver; John Peterson, Larimer County Hospital, Fort Collins.

Public Relations: Charles E. Levine, Chairman, Beth Israel Hospital, Denver; Ward Darley, M.D., University of Colorado Department of Medicine, Denver; A. Tergerson, Longmont Hospital and Clinic, Inc., Longmont.

SPECIAL COMMITTEES

Constitution and Rules: Owen Stubben, Chairman, Denver General Hospital, Denver; Harry Clark, Southwest Memorial Hospital, Cortes; Sister Mary Lina, St. Francis Hospital, Colorado Springs.

Hospital and Professional Relations: Roy Anderson, Chairman, Presbyterian Hospital, Denver; G. A. W. Currle, M.D., Colorado General Hospital, Denver; Louis Liswood, National Jewish Hospital, Denver; C. S. Biuemel, M.D., Mount Airy Sanliarium, Denver.

Rates and Charges: DeMoss Taliaferro, Chairman, Children's Hospital, Denver; Megr. John R. Mulroy, Catholic Hospital, Denver; Roy Frangley, St. Luke's Hospital, Denver; Elton A. Reces, Alamoa Community Hospital, Alamoas; Boy Anderson, Presbyterian Hospital, Denver; Richard Connor,

Resolutions: Sister Mary Raymond, Mercy Hospital, Denver; James A. Harrison, Community Hospital, Boulder.

Accuracy and Speed in Prescription Service DORR OPTICAL COMPANY

421 16th Street

Denver, Colorado

KEystone 5511

МІ

Cortone

Safety in the Prolonged Control of RHEUMATOID ARTHRITIS



Successful clinical experience with CORTONE in many large series of patients reveals the safety of this product. The administration of CORTONE does not necessitate any measures that are not readily available to the physician in everyday practice. The use of simple laboratory tests (sedimentation rate, urinalysis, blood count, blood pressure, and recordings of weight), individualized adjustment of dos-

age, and careful clinical observation will permit most patients to benefit materially . . . α without fear of undesired effects.

One investigator notes: "We have not been impressed by the severity or frequency of side-effects... The side-effects due to excessive adrenal cortical hormone disappeared when the hormonal agent was discontinued."

Norcross, B. M., N. Y. State J. Med. 51: 2356, Oct. 15, 1951.

CORTONE is the registered trade-mark of Merck & Co., Inc. for its brand of cortisone.



(CORTISONE Acetate Merck




MERCK & CO., INC.

Manufacturing Chamists

RAHWAY, NEW JERSEY
In Canada: MERCK & CO. Limited—Montreal

Latest Information on Penicillin Therapy



Ask Your Squibb Professional Service Representative

/OL

49

SS

2

5

2

IMI

... A New Squibb Aid for the Profession



Squibb, a leader in penicillin research and manufacture, presents the new edition of the Squibb Penicillin Handbook, "Important Principles Influencing Penicillin Therapy." It is based on most

recent clinical work and data of eminent authorities in the antibiotic field . . . new penicillin dosages . . . new recommendations for efficacy . . . oral and parenteral forms . . . combined therapy . . . drug resistance . . . therapeutic blood levels . . . reactions . . . continuous vs. discontinuous therapy . . . and many other subjects of interest to physicians.

Your Squibb Professional Service Representative will provide you with "Important Principles Influencing Penicillin Therapy" or any other Squibb visual and practical aids, without cost or obligation. Or you may write direct to E. R. Squibb & Sons, 745 Fifth Avenue, New York 22, New York.

SQUIBB A LEADER IN PENICILLIN RESEARCH AND MANUFACTURE





Long welcomed in home and institutional kitchens for its convenience, economy and flavor—frozen citrus is now acknowledged the "nutritive equal" of fresh. The Council on Foods and Nutrition of the American Medical Association has declared* that—under modern processing methods—approximately 98 percent of the vitamin C content can be retained in the frozen concentrated juice. And, when properly stored (below its freezing point), there is practically no loss of vitamin C. Frozen citrus can thus be confidently recommended for diets at all ages, including infancy.

*J.A.M.A. 146:35, 1951.

FLORIDA CITRUS COMMISSION . LAKELAND, FLORIDA

FLORIDA itrus

UMI

OL

Dr. Harris finds in difficult surgical accomplishments occur more frequently these days than when he first began his practice. His skill has continued to develop. Furthermore, improved anesthesia has helped make possible the regular performance of certain kinds of surgery which at one time would have been extremely risky, if not impossible linvestigators in academic laboratories have joined their efforts with those of the pharmaceutical industry for

Oh Doctor - I can really walk again Lilly



... the Development of Improved Anesthetics

"DOCTOR, your patient is ready." This classical statement as uttered by Morton in 1854, when he first demonstrated surgical anesthesia, is a phrase still commonly used by anesthetists. Since then, however, many developments have greatly improved anesthesia and aided the progress of surgery. Among these was the discovery of 'Metycaine Hydrochloride' (Piperocaine Hydrochloride, Lilly) in a university laboratory, where, in conjunction with Eli Lilly and Company, chemists were searching for a better local anesthetic. From a long series of chemically related substances which had been prepared, 'Metycaine Hydrochloride' was selected for extensive clinical evaluation. When used as recommended, it was shown to be no more toxic than procaine and to be capable of producing anesthesia more quickly, with greater certainty of effect, and for a longer period of time. These attributes of an improved local anesthetic are the reasons why 'Metycaine Hydrochloride' facilitates careful surgery.

Lilly

ELI LILLY AND COMPANY . INDIANAPOLIS 6, INDIANA, U. S. A.

UMI

5

2

OL

Montana New Mexico Utah Wyoming

Rocky Mountain **FEBRUARY**

Medical Journal

*Editorial

Russians Attend A Medical Meeting

THE Fourth International Congress on Mental Health was held in Mexico City in December, 1951. There were delegates to the Congress from thirty-two nations, including three psychiatrists from Russia. None of the Russians appeared on the published program but at the last plenary session a paper was presented by Dr. A. V. Snejnevsky, Professor of the Central Institute for the Perfecting of Physicians of the U.S.S.R. The paper was entitled "Principles of Prophylaxis of Psychic Diseases in the Soviet Union" and translations in English and Spanish were distributed among the audience before the paper was read. Dr. Snejnevsky read his paper in Russian while the interpreters read the prepared translations from their glassed-in booths at the back of the rostrum and the audience listened to the translations with head sets in the usual procedure of an international meeting. Unfortunately the paper was read quite rapidly and when the doctor finished. the interpreters were five minutes behind. hence a fair proportion of the audience who were not supplied with written translations heard the paper only in part. At the conclusion of the paper the chairman thanked the essayist in German but did not call for discussion.

The paper was remarkable in its content and here we quote the opening lines:

"In the Union of Soviet Socialist Republics, the country of victorious Socialism. the well-being and happiness of the people are the chief concern of the State. Care of the well-being of the population, of health of man is the law of the development of our State. 'Of all precious capitals of the world,' said J. V. Stalin, 'the most precious are people.' The socialist mode of production in the Soviet Union and the corresponding social order have done away with the laws of relative overpopulation inherent to capitalism which dooms millions of urban and rural workers to pauperism, beggary, diseases, and death."

Pavlov was featured as the hero of Soviet psychiatry and under his influence and that of Sechenov "Russian psychiatry has always tended to a physiological understanding of the nature of psychosis, and there lies its originally." Moreover Pavlov studied the social and family conditions of the patient and his conflict situations. Credit for the advance in Soviet psychiatry is also due to the peasants and the workers. "Under the guidance of science they have created in their socialistic State a firm base for a complete sanitation of the country, of working and living conditions for a successful development of the new system of socialistic health services. For the first time in the history of mankind the interests of the State fully correspond to the interests of the people."

The paper extolled the beauties of life under victorious Socialism in the new Russia. "The basic features of our society is general occupation in labor, the right to rest, guaranteed by milliard allocations for this purpose, maintenance in old age, sanitation labor and living conditions prevent diseases including nervous psychic illnesses."

When mental illness occurs among the Russians the patient receives medical care at the expense of the State, and as a rule the same doctor attends him in the hospital, in the outpatient department, and in his own home. Upon recovery the State finds suitable employment for him in a cooperative of invalids and if necessary the State provides a pension.

Many who listened to the paper thought that Dr. Snejnevsky was not altogether objective in handling his statistics. Not only had the incidence of dementia precox been reduced by half, but in the period between 1930 and 1948 the incidence of manic-depressive psychosis had been reduced to "six times less" while alcoholic psychosis had dropped to "ten times less." And currently there are no cases of alcoholic psychosis or general paresis in the Soviet Union. How accurate are these observations? Perhaps a little loose, for the doctor followed with the statement that "neurosis and psychopathy were not observed in the army and the rear during the Great Patriotic War." This appears contrary to human experience.

It was difficult to escape the impression that the doctor's thinking was politically polarized and that he was perhaps obliged to present propaganda rather than factual observations in medical science. Yet psychiatrists can be tolerant, and there is no doubt that those at the meeting shared the sentiments of the Director, Dr. John R. Rees of London, when he publicly welcomed "our friends of the Soviet Union" to the International Congress on Mental Health.

C. S. BLUEMEL.

REPORT OF COLORADO DELEGATES TO THE AMERICAN MEDICAL ASSOCIATION

The Fifth Annual Clinical Session of the American Medical Association was held in Los Angeles, December 4 to 7, 1951. Registration was, approximately, 4,500 physicians and about an equal number of guests. A summary of the activities of the Clinical Session and an abstract of proceedings of the House of Delegates may be found in the Journal A.M.A. of December 22, 1951, pages 1673 to 1702. This report is concise, comprehensive and excellently planned. As all members of the Colorado State Medical Society have access to the Journal A.M.A., your delegates believe that an extended report by them is unnecessary. However, it is urged that members read the report in the Journal. It is only 29 pages and can be read in one hour.

Particular attention is invited to: The address of President John W. Cline. The address of Dr. A. J. J. Rourke, president of the American Hospital Association, and of Mr. Donald Wilson, Commander of the American Legion.

"Guides for Conduct of Physicians in Relationship With Institutions."

The report of the Committee on Constitution and By-Laws.

The report of the Coordinating Committee.

The report of the American Medical Education Foundation.

Other items of interest: The Board of Trustees allocated one-half million dollars to the Education Fund for 1952. . . . The Society of Anesthesiologists presented \$2,500.00 to the Education Fund. . . . The House of Delegates authorized the Board of Trustees to pay "a liberal per diem allowance" to the President and Presidentelect in addition to expenses already authorized. ... The Board of Trustees was encouraged to "consider the purchase of suitable property in Washington, D. C., to serve as a permanent headquarters for the Washington office." . . . The Board of Trustees will report on the treatment of non-service-connected disabilities in Veterans' Hospitals at the June meeting in Chicago . . . The resolution of Dr. Eugene F. Hoffman of California on "Congressional investigation of the teaching of collectivism in schools" was ap-

At the Fourth Annual Public Relations Conference the address of Dr. Cyrus W. Anderson of Denver on "Explaining those 'other' medical expenses" was enthusiastically received. "Cy" did a splendid job both in the delivery of his address and in the question period. Mr. Harvey T. Sethman, who, in addition to being our own Executive Secretary, is now Chairman of the Advisory Committee to the Department of Public Relations of the A.M.A., presided at the afternoon session on December 3, 1951.

The program of the Coordinating Committee presenting United States Senators Taft and Byrd, both of whom made excellent speeches, was a great success. A Los Angeles businessman remarked that he was undecided whether Senator Taft was a good Democrat or Senator Byrd a good Republican.

At a dinner given to the House of Delegates by the Los Angeles County Society we were royally dined and were entertained by Mr. Edgar Bergen with Charlie and Mortimer, by Mr. Jean Hersholt and Hawaiian dancers.

Congratulations and appreciation are due the Los Angeles County Society, especially the cochairmen, Doctors J. W. Scott and L. Lafe Ludwig, Committee on Arrangements, for masterful and successful "staff" work.

Dr. A. C. Yoder, Goshen, Indiana, was elected General Practitioner of the Year.

> GEORGE A. UNFUG, WILLIAM H. HALLEY.

Original Articles

CRANIOCEREBRAL TRAUMA*

BASIC PRINCIPLES OF EXAMINATION AND CARE

ALEXANDER C. JOHNSON, M.D. GREAT FALLS, MONTANA

Increasing incidence of head injuries as a natural sequence of increasing mechanization of our daily lives is well recognized. It is obvious that the vast majority of such cases of injury to the central nervous system, and to its membranes and bony encasement, must come under the care of the family doctor. Excellent detailed works pertaining to trauma of the central nervous system in all of its aspects, notably those of Brock and of Rowbotham, are available. All such treatises suffer from the common fault of being primarily written by and for the neurosurgeon and neurologist. Such volumes obviously do not readily yield basic information to the family doctor and are liable to create a disproportionate perspective of essential and non-essential features in the management of these cases. The writer, whose work is in the field of neurologic surgery, has been impressed with the frequent deviation from sound physiologic concepts in the management of head injuries.

The most important single observation to be made when confronted with the patient who has incurred cranial and intracranial trauma is his state of consciousness. There is no more reliable index of the degree of generalized brain injury, though this does not preclude the possibility of serious local injury as may be associated with comminuted depressed fractures, where the general concussive effects are often reduced by the localized fracture of the skull. His state of consciousness, as noted initially, then gives a base line from which one may estimate evidence of returning consciousness or deepening of coma.

It should not be forgotten that the patient may not have merely skull and brain injury, but that he may incur distant traumatic lesions simultaneously; and, therefore, the general condition of the patient, particularly the vital signs and evidence of shock, should be evaluated. In this latter regard it should be stressed that true shock is not a characteristic part of craniocerebral trauma, and when present can rarely be explained on the basis of the head injury. A shock-like condition has been described with some brain stem injuries; but in the author's experience surgical shock, when present in a head injury case, has nearly always been due to other injuries present. This feature is of vital importance and all too often serious visceral or other injuries are overlooked because of the assumption that the state of shock is due to the head injury. As a matter of fact, serious brain injuries are associated with physiologic changes which are actually the reverse of the picture of shock, the patient having a full, slow pulse, elevated blood pressure, deep stertorous respiration, and hot, flushed skin, in contradistinction to the well-known features of surgical shock. It follows then that a brief, but careful, examination of the chest, abdomen, spine, and long bones is indicated.

The really essentially features of the neurologic examination of significance in the initial evaluation of the head injury patient are unfortunately often either omitted or excluded in favor of a detailed elicitation of various reflexes. It goes without saying that the patient who has regained consciousness should be the subject of a full and detailed neurologic examination, but let us deal primarily with the unconscious patient. We have already observed his general condition and state of consciousness.

^{*}Presented at a symposium on trauma at the U. S. Administration Hospital, Fort Harrison, Montana, April 19, 1951.

To be presented at the Scientific Session of the Interim meeting of the Montana State Medical Association, February 29, 1952.

During this time we have probably noted the presence or absence of any gross paralysis. Does the patient move about in bed and does he move both sides equally? If he does not move spontaneously does he respond to painful stimuli such as pinching the arm or leg? Facial paralysis can often be detected by pressure over the supraorbital nerve, producing grimacing. These reponses to painful stimuli also serve as an index of the profundity of coma in patients in whom there is no spontaneous movement. The head and cranial orifices should be examined, noting particularly the presence of blood in the external ear canals, which practically always indicates a petrous pyramid fracture, and the presence of any scalp lacerations or other evidence of local trauma. The spine should be inspected and palpated for gross deformity and due caution should be given always to the possibility of spine fracture in association with head injury. This is particularly noted in the so-called "Coleman's Syndrome" in which skeletal damage to the shoulder girdle, associated with head injury, is strongly suspicious of the possibility of a cervical spine injury.

The eyes should be examined for conjugate alignment and the size and equality of the pupils. Examination of the fundi is not necessary immediately as funduscopic changes are not to be expected initially, except in those cases of massive and rapidly fatal intracranial hemorrhage. Under no circumstances, and this should be emphasized, should any sort of mydriatic drug be used in examining the fundi of head injury cases, since this is not only unnecessary but destroys some of the most valuable neurologic findings. It is indeed the rare fundus that cannot be adequately examined without the use of mydriatics and in these rare cases it is preferable to preserve the pupillary responses rather than to examine the fundi.

A comparison of the common muscle stretch reflexes, the biceps and triceps in the upper extremity and the knee and ankle jerks in the lower extremity should be made and recorded. The plantar reflex should likewise be noted as to the presence or absence of the Babinski sign.

It is not unintended that the consideration of x-ray films of the spine and skull has been left to the last. To paraphrase the late and great Dr. Harvey Cushing, it is apparent that too much attention is paid to disturbance of bony continuity and too little to the disturbances of central nervous system physiology. This faulty thinking is emphasized in the common practice of referring to head injury as synonymous with skull fracture; while, as a matter of fact, there are no symptoms directly referrable to a fractured skull but only to the associated trauma to the underlying brain. With this in mind, it is then apparent that x-ray films are made only when the condition of the patient warrants, and under no circumstances should a critically injured patient be moved about for the purposes of obtaining x-ray films, which will have no bearing on the immediate management of his problem. When his condition has stabilized sufficiently to warrant x-ray films these should be made, using the best possible technic and obtaining at least the P. A. and Towne (occipital) positions and both lateral views. Films of poor diagnostic quality or an inadequate number of views to properly evaluate the pathology that might be disclosed are of little help.

Let us now turn to the initial orders that might be written in a case of serious craniocerebral injury, and the rationale of each. The blood pressure, pulse, and respiration should be determined at half-hourly intervals initially and recorded on a graph in order to follow changes in the vital signs. Let it suffice to say that a progressive slowing of respiratory and pulse rate, particularly associated with rising blood pressure and a rise in temperature, is classically characteristic of increasing intracranial pressure. These changes may not all be evident simultaneously. The nurses should be well instructed as to these facts, as uninformed nurses may assume that as long as the pulse is "strong" and the blood pressure "good," the patient is in satisfactory condition. His state of consciousness should be observed and the nurses likewise instructed to report immediately any progressive changes in vital signs and state of consciousness, or the developing of inequality of the pupils, as well as the occurrence of vomiting or convulsions, all of which may be strongly indicative of an increase in intracranial pressure, as associated with certain surgical complications to be discussed subsequently. Too often the casual notation "resting quietly" in the nurses' notes actually indicates developing and unrecognized coma.

The head of the bed is elevated one-third. which decreases the intracranial venous pressure, and accordingly decreases the cerebrospinal fluid pressure. This also provides a greater pulmonary vital capacity. The patient should be turned completely from side to side at least every two hours and thoroughly suctioned, using a catheter of about No. 14 size with multiple holes in the end, and passing this catheter far down through each nostril into the region of the larynx and hypopharynx. If this stimulates coughing so much the better. After the first few hours the possibility of aggravating any intracranial hemorrhage in this manner is academic; and if an epidural or subdural hematoma is to develop, the groundwork has already been laid before this time. This writer is continually amazed that these simple and fundamental procedures in the care of an unconscious patient seem to be generally untaught in schools of nursing. A small pillow under one shoulder is thought to pass for turning the patient, and the removal of a small amount of saliva from the anterior part of the mouth with a tonsillectomy suction tip is little better than complete neglect. Where secretions are profuse postural drainage for fifteen or twenty minutes prior to suctioning is useful. Atropine is of too short duration to be really useful in this respect. In rare instances tracheotomy must be done. It is probably true that with the exception of unrecognized surgical complications, such as chronic subdural hematoma or brain abscess, most late fatalities from craniocerebral trauma are due to pulmonary complications arising from faulty nursing care.

In spite of the fact that every medical corpsman in the armed forces during the recent World War was thoroughly instructed not to use his morphine syrette in any casualty in which a head injury was present, morphine is still almost invariably and unfortunately used in civilian cases, at times even in semi-conscious or unconscious patients. At no time should morphine or any of the opiates be used because of the depression of respiration, which is often already severely compromised in head injury cases. The resultant increase in intracranial venous and cerebrospinal fluid pressure resulting from such respiratory depression, and the interference with pupillary light reflexes are dangerous secondary effects to be avoided. The only exception is the occasional case having severely painful injuries elsewhere, in which small doses of Demerol combined with barbiturate may be judiciously used, but never to the point of significant hypnosis or narcosis. Where possible, no sedation whatsoever is best, but in the occasional extremely restless patient small doses of sodium phenobarbital may be used intramuscularly, but again, never enough to produce somnolence. At all times it must be remembered that the state of consciousness is a primary index of the patient's condition, and nurses should be informed in this regard so that developing coma is not simply recorded in the nurses' notes as "sleeping quietly." If necessary bed rails and restraints of the extremities may be used where there is marked restlessness. It should be kept in mind that restlessness often indicates bladder distension in a patient not able to make his wants known. Catheterization is a safe "sedative" in such instances.

The question often arises as to the appropriate fluid intake in patients unconscious following craniocerebral trauma. It was once quite in vogue to drastically reduce fluid intake and even to administer purgatives, as well as quantities of hypertonic fluids by vein. It is often noted that where patients are subjected to such drastic dehydrating measures they may not only fail to show the desired improvement, but rather their condition may progressively worsen. The writer has seen such cases show immediate favorable changes following administration of restorative quantities

of fluids intravenously, resulting in a prompt fall in temperature, "lightening" state of consciousness, and generally improved appearance. It is this writer's opinion, and probably one shared by the majority of neurosurgeons, that an adequate fluid intake, on the order of 2,000 to 2,500 c.c. per twenty-four hours, is indicated; more in dehydrated or febrile cases. Patients under the writer's care receive this amount intravenously for the first two or three days (appropriate volumes in children), following which a Levine tube is inserted through the nose if it appears that prolonged unconsciousness is to be expected. Through this stomach tube the appropriate total volume of fluid is given, including a high vitamin liquid diet of about 2,000 calories, or higher if the temperature has been elevated. It is not felt that the use of hypertonic intravenous fluids is generally justified, except where additional time to prepare for emergency surgery in an acute compression syndrome (epidural or subdural clot) is necessary. This may shrink the brain sufficiently to postpone slightly the exact time of irreversible compression. The use of hypertonic fluids as part of routine treatment not only does not have quantitatively as much effect as generally thought; but furthermore, it has been well shown that the clinical diagnosis of posttraumatic cerebral edema does not necessarily coincide with this actual physiologic state, and therefore, hypertonic fluids might be administered in a case in which cerebral swelling is not the pathologic mechanism concerned. Since this writer only uses hypertonic fluids as a preoperative emergency measure, hypertonic (50 per cent) dextrose is the solution generally used. As surgical relief is anticipated the well-known secondary rise in cerebrospinal fluid pressure is of no moment. Under other circumstances hypertonic plasma or serum albumin may be preferable.

Hyperthermia may occur as a complication of brain injury in any degree from a very slight elevation of temperature to a progressive hyperpyrexia with fatal termination. Apparently the presence of blood in the cerebrospinal fluid may act as an irritant or otherwise as a disturbing influence on the thermal regulation of the body to produce of itself some degree of elevation of body temperature. With higher degrees of body temperature, and particularly the severe cases progressing to a fatal hyperpyrexia, the presence of significant damage to the heat regulatory centers of the hypothalamus may be presumed. This fact is well known from the occasional occurrence of hyperpyrexia following surgical intervention for tumors involving the region of the diencephalon, particularly suprasellar and third ventricular tumors. Generally speaking, the development of hyperthermia due to disturbance of central nervous regulation will be noted soon after the occurrence of trauma and any elevation of temperature first noted as long as twelve or eighteen hours after trauma should be regarded critically as likely due to other causes, particularly pulmonary complications. The treatment of the hyperthermia itself consists of measures aimed at increasing the loss of body heat and, in addition, measures to resist the catabolic effects of the elevated body temperature.

Covering the patient's body with damp towels seems to be the most effective method of lowering body temperature since the evaporation of this moisture requires a tremendous amount of heat. It is not necessary and in fact undesirable that the towels be cold as it is the evaporation of the water and not its temperature that is responsible for the loss of body heat effected. This is most effective where the skin temperature is elevated. Where there is a cutaneous vasoconstriction with a cold skin surface the application of these damp towels should be preceded by a brisk rub of the skin with tepid water and rough sponges or towels to induce vasodilatation. The effectiveness of this treatment may be increased by directing electric fans over the coverings of the patient to increase evaporation. Alcohol sponges are also useful and may be combined with brisk massage of the skin to increase cutaneous circulation. The writer is unimpressed with the effectiveness of ice bags since these usually produce a rather prompt cutaneous vasoconstriction with increased retention of heat, sometimes greater than the amount lost in melting the ice by conduction.

The catabolic effects of hyperthermia are combatted by the use of an oxygen tent, primarily to supply the additional oxygen needed because of the elevated body metabolism, but also for some cooling effect by conduction, if the temperature of the tent is maintained at proper levels. The fluid intake should be increased sufficiently to keep urinary output and urine concentration within normal limits, and supplementary intravenous fluids or high-calorie tube feedings are desirable to supply additional caloric intake. These measures are useful in treating any excessive hyperthermia; though some cases of severe and irreversible hypothalamic damage will be found to respond little, if at all, to these or any other measures and generally terminate fatally.

A word should also be mentioned in regard to the use of lumbar puncture in head injuries. As this writer has pointed out in another article, the cerebrospinal fluid findings bear little relationship to the clinical management of the head injury patient. It is well known that while the cerebrospinal pressure is generally elevated following head injury it may be normal or low even in severe cases. Furthermore, the presence of a high spinal fluid pressure does not necessarily indicate the presence of an intracranial surgical lesion and occasionally a surgical lesion, particularly a chronic subdural hematoma, may be present without remarkable or even no elevation in spinal fluid pressure. The fluid itself may be clear to grossly bloody, and the degree of extravasation of blood into the cerebrospinal fluid likewise bears no direct relationship either to the seriousness of the injury nor to the proper management of the case. It should be again emphasized that the injudicious use of spinal puncture in the presence of an expanding intracranial lesion carries just as much, if not more, risk in the case of a post-traumatic hematoma as in a brain tumor. Greater adjustment of the brain to the more slowly expanding tumor is possible. Lumbar puncture in such cases may not only produce a rapid fatality in

some instances, due to herniation of the cerebellar tonsils through the foramen magnum and the transtentorial herniation of the temporal lobe through the tentorial incisura compressing the brain stem; possibly more often the puncture produces only an accelleration of this train of events converting a favorable operable case, in a relatively short time, into one beyond surgical reversibility. While there is no entire agreement among neurosurgeons on the value of diagnostic lumbar puncture in head injuries, it seems to be rather well agreed that this procedure is of no therapeutic value in the form of repeated spinal drainage, as once advocated widely. The writer finds the principal value of lumbar puncture to be in those cases admitted to the hospital unconscious without adequate history, in which case lumbar puncture is only a part of the various studies, including blood sugar, n.p.n., etc., indicated in the differential diagnosis of unconsciousness.

Let us direct our attention now to the consideration of surgical principles. First, it should be realized that any person who has incurred a head injury, even an apparently minor injury, may develop a complication of urgent and grave surgical implications: therefore, such possibility must be considered in any head injury case showing progressive neurologic impairment and development of stupor or other evidence of increased intracranial pressure. One of the earliest signs may be the development of incontinence in a patient who previously was sufficiently rational to ask for the bed pan or urinal. Delay in surgery may result in the development of irreversible cerebral compression with fatality even after surgical evacuation of the hematoma; therefore, the suspicion of increasing intracranial pressure is adequate reason for immediate neurosurgical evaluation and intervention if indicated. There is no more distressing situation to the neurosurgeon than to receive a patient in decerebrate rigidity with terminal and irreversible cerebral compression as the result of a completely curable condition, such as a subdural or epidural clot, as a consequence of excessive delay for "observation" in the attempt to arrive at a positive diagnosis. Particularly in the case of the epidural hematomas, which being generally of arterial (middle meningeal) origin are rapidly developing, the entire course of events from injury to fatal termination may be but very few hours. The mortality from this lesion today approaches if not exceeds 50 per cent, in large part from delay in surgery. Dramatic recoveries are to be expected in most cases where surgical intervention is undertaken early, provided there is not severe associated brain injury.

Compound skull fractures necessitate prompt surgical intervention for the same reason as compound fractures elsewhere, primarily to prevent infection. In addition, debridement of the underlying brain and meninges and the repair of the meninges is indicated to prevent infection and minimize cortical scar formation. This, however, does not apply to compound linear fractures except where foreign matter, such as hair, may be trapped in the suture line. The "allor-none" rule should apply, and no superficial preliminary debridement or repair of the wound should be undertaken as this may only promote spread of latent infection. The entire procedure should be done at one session by one trained and equipped for major brain surgery, as such injuries are often more serious than the appearance of the x-ray films or the patient's condition might suggest.

Missile wounds, for practical purposes, may be considered as compound craniocerebral injuries to be managed with the same principles as other wounds of this groupessentially, complete definitive debridement, repair, and arrest of hemorrhage, as soon as the patient's condition permits. There are other considerations not within the scope of this paper. An operative mortality rate on the order of only 14 per cent from all theaters in World War II for such cases reaching neurosurgical units attests to the soundness of such management. While missile wounds of the brain are not common in civilian life, the writer believes the mortality rate in such cases is generally higher because of the erroneous conception of such wounds as "always fatal."

Cerebrospinal fluid fistulas may occur from the nose, the ears, or directly through a compound fracture. Those occurring through the nose are of the greatest direct significance since a spinal fluid leak occurring through a compound fracture will be simultaneously corrected with surgical repair of the fracture itself and the underlying meninges; while cerebrospinal fluid otorrhea is generally self-limiting, presumably because of the distance that this fluid must travel through the fractured petrous pyramid. Practically all cases of cerebrospinal otorrhea will stop spontaneously within a few days at the most and surgical intervention is but rarely required. The patient should be on intensive prophylactic chemotherapy, preferably of the sulfonamide drugs because of the greater levels obtained in the normal cerebrospinal fluid circulation. Cerebrospinal fluid rhinorrhea presents quite another problem. While many such cases do stop spontaneously, and initially are managed similarly to cases of cerebrospinal fluid otorrhea, the lack of any great bony barrier greatly facilitates the continuation of such a leak, since these leaks ordinarily occur through fractures into the frontal sinus or inferiorly into the cribriform plate and ethmoids. The patient should be forbidden from blowing his nose, coughing, and straining, since these actions increase the leak of fluid. A real and serious danger of meningitis, often fatal because of its mixed bacterial nature, is present in such cases. Prior to chemotherapy and the antibiotics it was desirable in such cases to wait not more than a few days before surgical intervention. At the present time even with the aid of our potent antibacterial agents it seems highly dangerous to wait more than a week before considering surgical intervention with repair of the meningeal laceration and its fistulous connection. The presence of intracerebral air is pathognomonic of this condition even where there has been no recognizable cerebrospinal fluid leak, which may be intermittent and of a degree not recognized by the patient.

A word should be mentioned in regard to exploratory burn openings or "inspection holes," as termed by Rowbotham. There are many cases in which evidence of serious generalized brain injury makes the evaluation of a developing surgical lesion difficult, and in such cases rather than delay further, exploratory trephine openings may be made under local anesthesia (a procedure of ten or fifteen minutes time) with no detriment to the patient in the event of negative findings. Such a procedure may result in early removal of a not clearly recognized hematoma. The safety of this procedure should be emphasized since all too often the patient's family and physician consent to such surgery with misgivings or dangerous delay only because of the erroneous idea of long, serious, and hazardous surgery.

A question which often arises here in the West, where distances to neurosurgical attention are great, is the matter of transportation. The experience of the recent war conclusively showed that head injury cases may be transported quite well, provided there are not severe associated injuries with resultant shock. After shock is treated in the usual manner and the vital signs have become stabilized, these patients can then be moved readily. Whether transported by airplane or ambulance, nurse attendance and the availability of oxygen is highly desirable, the latter particularly in patients being transported by air. It should be noted in this regard that time is of the essence and the patient, whose life may be saved by prompt surgery, will be no better for delay and indeed may reach such moribund condition that transportation is out of the question. When a patient has reached such a state surgery will usually be of no avail. As a general rule it may be stated that the patient with a neurosurgical emergency will get no better by delay, and if suitable for surgery at all he is generally also suitable for transportation. The writer believes that such a patient, not yet in moribund condition, can be moved with relative safety. It is not to the patient's advantage that the neurosurgeon be brought to the patient to attempt major neurosurgical intervention in an unfamiliar hospital with an untrained operating team. The time lost not only will be greater and the surgery more hazardous but, in addition, the pa-

tient must be then deprived of the postoperative observation of the neurosurgeon.

Summary

The management of craniocerebral trauma cases consists of the initial evaluation of the patient, including his general condition, the presence of other injuries in addition to the head injury, the treatment of shock if present (and the recognition that this is nearly always not due to the head injury, but must be explained by other injuries to be sought for in the examination), and a brief neurologic evaluation to include evidence of paralysis, head and cranial orifices, ocular position and pupillary size and equality, and the equality of the reflexes and presence of pathologic reflexes. The follow-up and care then consists of a careful evaluation of the state of consciousness, neurologic and vital signs at frequent intervals, maintenance of normal pulmonary aeriation and prevention of pulmonary complications, and prompt neurosurgical attention where indicated.

NEW TYPE OF INTERNSHIP

Beginning July 1, 1952, a new type of internship will be offered in Colorado, sponsored by the participating private Denver hospitals and the University of Colorado School of Medicine. The new Community Hospitals Internship will be tried for one year and if it proves successful, it may become a regular part of the training of young physicians in this area. It will be in addition to the regular Denver General Hospital internship which has been offered for many years and which will remain essentially unchanged.

In the new internship, the young physician in training will spend six months at Denver General Hospital and six months at one of the private participating hospitals in Denver. It is expected that this plan will better utilize the unique teaching material available in private hospitals, will offer the intern a broader exposure to different viewpoints and technics in the practice of medicine, and will provide a better distribution of internship services to meet the community needs for both private and indigent patients. It will offer the young physician an opportunity for experience in private hospital practice.

In addition to the usual ward assignments, the intern will also spend part time in the newly established General Practice Clinic during his six months at Denver General Hospital.

The new Community Hospitals Internship has been approved by the Council on Medical Education and Hospitals of the American Medical Association and also by the National Interassociation Committee on Internships. This is the first instance in this country of an internship combining equal experience in private and public hospitals. If successful, it may well become a pattern for internships in other parts of the country.

MEDICINE'S OPPORTUNITY OF A LIFETIME*

ERVIN A. HINDS, M.D.

We were just getting back to living like human beings again following World War II when right out of the clear, the precursors of World War III poke their ugly heads. We have been caught frustrated, angry, confused, deeply disappointed and badly off balance. Our national thinking got snarled up with uncertainty and alarm. Like frightened animals many people started biting at each other. But it seems to me that we, as doctors, are living in one of the most exciting and interesting periods of medical history-and I don't mean medical history from only the viewpoint of scientific achievement. I mean that this is the most interesting time in all history from at least three viewpoints-scientific, economic and political.

We are thankful that medicine as a profession abandoned the sidelines right after World War II and got into all three of these activities, whereas before it had taken part in only one—the parade of scientific progress. Now that medicine has limbered up its economic and political muscles, it has really come of age in civic affairs—and that is why I say that medicine has an opportunity of a lifetime, particularly in the next few years.

It was not long ago that Dr. Elmer Henderson, then President of the A.M.A., observed: "American medicine has come a long way in a short time." He was referring primarily to the economic and political awareness that doctors and medical societies have developed in the period since Oscar Ewing's now notorious meeting in Washington in 1948 which he called the National Health Assembly. You will recall that his so-called National Health Assembly had been framed in advance, which made the recent basketball fixers look like pikers. When his assembly adjourned, it had failed to deliver medicine into Oscar Ewing's hands as he had planned. So he and his cohorts re-wrote and re-worded the supposed report of the conference which, you will recall, also declared war against medical free enterprise.

Medicine accepted the challenge and all of us, through our American Medical Association, started a national educational campaign to carry our political battle right to the people. Most of the doctors in the country took part personally. Most county medical societies worked hard. Almost every state medical society, certainly including our own—did excellent jobs. So did many allied organizations and individuals. And in this three-year period, medicine has found many friends which it did not know it had.

One of the turning points in this war was the 1950 general election. And when I say "war," I mean exactly that, because it has been, it is, and it will continue to be a war of survival for free enterprise. Many aspects other than the medical aspects of this war came to a focus at the 1950 general election which, as we now all know, resulted in a great victory for those who believe in free enterprise. This activity has given us all a prominence in civic affairs that we have never had before. This again points up my theme that medicine has, right now, its opportunity of a lifetime. Our recently acquired civic and political prominence gives medicine the opportunity to recapture that leadership. Certainly these opportunities include leadership in our communities, in our community political affairs, in the conduct of our civic organizations-in fact, in every aspect of modern

If we want to make the most of these opportunities, we must go on with renewed vigor, not only in support of our scientific and professional standards, not only in support of our belief in free enterprise throughout American life, but in support of the hundreds of projects that arise in our daily life that help the average citizen of our country to a high standard of living and a better appreciation of America. It means more devotion to our medical societies and our civic enterprises—it means

^{*}Read at the Eleventh Annual Western Colorado Springs Clinic, Grand Junction, Colorado, April 1, 1951.

that every doctor must devote a little more time to his professional community and to his civic community than he has done in the past. Putting it another way, it means devoting more interest to our fellowman and a little less to ourselves.

Freedom is the most important heritage we have. It always presupposes, however, a personal acceptance of responsibility. If a people will not assume this individual responsibility they will eventually lose their freedom. Anyone who selfishly violates the moral law in the field of business, the professions or labor, thereby threatens our freedom. Man must either control himself from within or eventually he will be controlled from without. To remain free, we must be responsible men.

You see, if I have made myself clear, I am trying to explain to you why I believe medicine, the individual doctor, the county society, the state society, and the A.M.A.medicine as a whole-has the opportunity within the next two or three years to lead the thinking of this nation of ours. If we accept this opportunity and exploit it properly, medicine may change the history of our nation at the general election in 1952. It is my belief that on that date we will either start the long road back to the kind of government and kind of free enterprise that made this country great, or else we will slip further into the quicksands of socialism.

To grasp this opportunity we must have clean hands. We must do more than render lip service to our ethics and our moral standards. In plain language, what I mean is that any doctor who is guilty of overcharging, of overtreatment, or of neglect of patients must be controlled, disciplined and, if necessary, must be removed from our midst. We must give the best medical care of which we are capable to all the people, and we must do so at a cost which each person can afford to pay. We cannot wait about cleaning our own house. We have made a good start, but we must move faster because every time any doctor imposes upon

a patient, another voter is ready to exchange free enterprise for the sugar-coated nostrum of a bureaucrat. Remember that with the critical international situation as it is, the bureaucrats are intensifying their efforts to get control of all free enterprise, including medicine, by labeling every one of their proposals as "emergency measures in the interests of national defense." Thus we have no time to waste.

Not only must we have clean hands, but we must all pull together. Up to now, medicine has always kept its organization in a most democratic manner, and it should ever remain so. Those like myself, however, who have found themselves temporarily in a position of leadership, wish that in developing these opportunities, more doctors would take part in the preliminary discussions, in the meetings of their county medical societies, in discussions with their delegates, in meetings of committees and boards. If they would express their ideas before decisions are made, decisions probably would be wiser; however, once committees and boards have made their recommendations, and houses of delegates have considered them and arrived at decisions, policies thus made should be considered binding upon all of us. Let's all take part in making our medical policy decisions. Then let's really pull together and gain that leadership I have tried to describe to you.

Every one of us can start on this road in our own offices, when we see our first patient tomorrow. Every one of us can keep everlastingly at it, bearing in mind that we honestly deserve the opportunity of a lifetime to help keep our nation great. What every one of us does from today forward will either increase or lessen that opportunity. Each of us has the confidence of a sufficient number of people, and, fortunately, we have strong medical organizations, local, state and national, to carry on that part of it. A few years ago, you pulled off your coats and doubled up your fists and decided to make a fight of it. The first two or three rounds are over with and we are stronger than when we began. Now, let's finish the job.

PREVENTION OF POSTOPERATIVE PULMONARY INFECTIONS*

BY INHALATION OF MI CRONIZED PENICILLIN

JOHN A. DIXON, M.D., SALT LAKE CITY, UTAH

and

EARLE B. MAHONEY, M.D.

ROCHESTER, NEW YORK

Pulmonary infection is still a fairly common postoperative complication in spite of advances in our knowledge of anesthesia and postoperative care. The use of antibiotic agents such as penicillin has reduced the seriousness and mortality of pulmonary infection but considerable morbidity is still attributable to this complication. The use of penicillin in a micropulverized form administered by inhalation pre-operatively has been reported in a previous article by Taplin, Cohen and Mahoney. This preliminary investigation indicated that postoperative pulmonary infection was greatly reduced by prophylactic penicillin administered by inhalation. Intramuscular penicillin did not seem to exert a comparable beneficial effect. The purpose of this report is to summarize recent experience and to further amplify the rationale for the use of micronized penicillin in the light of recent experimental data.

Methods and Materials

Using the type of operation as the sole criterion for selection, 150 cases were chosen for study at the Strong Memorial and Rochester Municipal Hospitals during the period July, 1949, to February, 1950. A deliberate attempt was made to select those cases wherein postoperative pulmonary infections were most likely to occur, i.e., upper abdominal cases such as vagotomies, cholecystectomies and gastrectomies, and thoracic cases such as thoracotomies, lobectomies, and pneumonectomies. Included in this group were seventy-seven males and seventy-three females, the average age being 50.8 years.

The day before operation, a nasopharyngeal culture was taken, plated for penicillin sensitivity and 200,000 units of micronized

penicillin given by inhalation. The day of operation, 100,000 units were given before operation and 100,000 units after operation when the patient had reacted sufficiently to cooperate; 100,000 units were given each day for the first two postoperative days and on the third day, another nasopharyngeal culture was taken and the penicillin sensitivity of the flora again determined.

The distribution of the cases according to type of operation appears in Table 1.

TABLE 1

Type of Operation and Distribution of Pulmonary Complications

Type of Operation—	Number of Cases	Complica- tions
Gastrectomy	27	3
Cholecystectomy		1
Transthoracic sympathectomy	y 30	1
Colectomy	12	0
Thoracotomy	10	1
Resection sigmoid	9	0
Vagotomy	8	0
Pneumonectomy		1
Lobectomy	5	1
Exploration common duct	4	0
Exploratory laparotomy	4	0
Lumbar sympathectomy	3	0
Abdominoperineal resection		0
Hernia	2	0

One case of splenectomy, diaphragmatic hernia, nephrectomy, Meckel's diverticulum, vulvectomy, ventral herniorrhaphy and thyroidectomy.

In classifying postoperative pulmonary infections, three groups were recognized. First, the patients who had a cough, occasional scattered moist râles, a mild temperature elevation and no x-ray evidence of atelectasis or consolidation, were said to have bronchitis. Second, those patients who were found to have a moderate elevation in pulse or respiratory rate and on physical examination were found to have a shift of the trachea, either absent or broncho-vesicular breath sounds with râles, mild

^{*}From the Department of Surgery, University of Rochester School of Medicine and Dentistry, Rochester, New York, and the Surgical Services of Strong Memorial and Rochester Municipal Hospitals.

fever and x-ray evidence of atelectasis were classified as having atelectasis. Third, those patients who had any or all of the above signs and symptoms plus a fever and x-ray evidence of an inflammatory infiltration were classified as having broncho-pneumonia.

Results

In the 150 cases studied there were eight cases in whom postoperative pulmonary infections were encountered. Of these cases there were seven patients who had atelectasis and one case of bronchopneumonia. This is a complication rate of 5.3 per cent for the series. Contrary to the findings of other investigators, there were almost twice as many males as females who had infections, the relative percentages being 6.8 per cent and 3.9 per cent, respectively. The average age of the group with complications was 46 years. These results appear in Table 2 along with fifty-one previous cases reported from this hospital. Included for comparison are three series of control groups selected at random from previous years and forty cases given aerosol penicillin prophylactically. In all, 236 control cases showed a remarkably constant incidence of postoperative respiratory infection from year to year averaging 19 per cent; 201 cases treated prophylactically with micronized penicillin had a complication rate of 4.5 per cent. The control series were selected insofar as possible to include comparable operative, seasonal, and age groups.

The results of the bacteriological study appear in Table 3. The high degree of penicillin sensitivity of the common organisms of the nose and throat as well as their disappearance following micronized penicillin is evident. The low incidence of pneumococci is probably a result of the bacterial technics utilized rather than being representative of the true carrier incidence of the population, which according to some investigators may be as high as 50 per cent. In determining the latter figure, serial cultures, throat washings, and reculturing of numerous colonies was done, which undoubtedly would uncover a number of isolated colonies of pneumococci that would be overgrown or missed by the usual methods. It is interesting that the case in this series from which pneumococci were isolated was the only case of bronchopneumonia encountered.

In the administration of any drug to a large number of persons the question of sensitization always arises. Careful examination of the patients given micronized penicillin was carried out for evidence of sensitization and no instance of urticaria, pruritis, dermatitis, arthralgia, rhinitis, or asthma which could be attributed to the drug was found. In three cases, or 2 per cent, a mild stomatitis was noted, consisting of superficial small mucosal ulcers over the buccal and pharyngeal mucous membranes, which in no case necessitated discontinuance of the penicillin and in every case disappeared rapidly following cessation of

TABLE 2 Incidence of Postoperative Pulmonary Complications

No. of Cases			Postoperative Pulmonary			Complications	
		Pre- or Postoperative Micronized Penicillin	Bronchitis	Broncho- pneumonia	Atelectasis Without Pneumonia	Total	No. Complica- es tions Pet.
50	Feb., March '47	None	- 2	4	3	9	18
50	June, July '47	None	1	7	2	10	20
96	Nov. '47-April '48	None	3	11	3	18	18.7
40	April, May '48	Atomizer or Aerosol Penicillin	0	8	0	8	20
51	May, July '48	Micronized Penicillin	1	0	0	1	2
150	July '49-Feb. '50	Micronized Penicillin	0	1	7	9	5.3
O	over-all Averages	(1948-1950).					Per Cent Complications
Witho	out micronized pe	nicillin	236		*************************		19
		cillin				100000000000000000000000000000000000000	4.5

therapy. Eight cases having two-stage procedures, such as sympathectomy, received micronized penicillin twice at intervals of from one to six months and in none of these was a reaction noted during the second period of administration.

Though probably not on a sensitivity basis, 6 per cent of cases reported transient nausea while inhaling the penicillin. It is interesting that all cases of nausea and stomatitis occurred in non-smokers who represented a relatively small percentage of the total number of cases.

Discussion

Comparing and evaluating statistics on postoperative pulmonary infections is always difficult, due to the variations in "respiratory awareness," and the technics of observation and classification. Various investigators have reported series with complication rates from 2 to 69 per cent. King emphasizes that the greater the interest in this condition and the more it is studied, the greater will be the number of cases diagnosed. In this series of cases an attempt was made to eliminate as many variables as possible by having the same individual check the pulmonary status of all patients pre- and postoperatively and record the complications on the basis of his findings. All cases of infection were proved by positive x-ray findings, as were suspicious cases eliminated by negative x-rays.

That the basis of postoperative pulmonary infections is atelectasis has been shown by many investigators, notably Coryllos and Birnbaum and later Mousel. The pathogenesis of atelectasis is described as follows: A mucous plug obstructs a bronchus, the air in the lobe supplied by that bronchus is rapidly absorbed, and collapse occurs. Bacteria in the trapped secretions rapidly invade the collapsed segment and bronchopneumonia follows. Recent studies by Baarsma and Dirken on collateral ventilation have shed much light on the mechanism of such collapse, which is applicable in arriving at a rational basis for the prevention of postoperative atelectasis and pulmonary infections. These investigators found by experimental studies in rabbits

and clinical observations in man that occulsion of the main bronchus to a lobe regularly produced atelectasis. Occlusion of a bronchus beyond the first primary division of the main bronchus to a lobe did not produce atelectasis in healthy subjects. The reason given for this phenomena is that collateral ventilation from the surrounding lobules which are supplied by unobstructed branches of the primary main bronchus occurs via the interalveolar pores of Kohn, thus preventing collapse. It was found that this collateral ventilation may amount to as much as 60 per cent of the volume normally supplied by the bronchial ramus to the pulmonary lobule. That these pores were not fissures of tissue or artifacts was demonstrated by special fixation technics in which the pores were visualized and by inference in that the permeability of such communications did not increase with increasing pressure and were open at even the most minute pressure differences. Most important to the problem at hand is that it was found with occulsion of a secondary bronchus, that atelectasis occurred only when collateral ventilation was eliminated (1) by shallow breathing, or (2) by the development of an inflammatory process in the unobstructed portion of the lobe. In the first case the communications or pores are obliterated by simple collapse and in the second case by the occurrence of in-

TABLE 3					
	No. Cases Pre-Op- eratively	Sensi- tive Pet.	Present Postop- eratively		
Strep. viridans	88	100	5		
N. catarrhalis	49	96	5		
Diptheriods	30	100	3		
S. albus hemolyticus	31	100	4		
S. aureus hemolyticus.	20	95	4		
Parainfluenzae hemolyticus	16	100	1		
Yeast		0	21		
B. aerogenes	8	0	33.		
E. coli	10	0	47		
B. proteus	2	0	4		
Non-hemolytic strep	2	100	0		
Strep. hemolyticus	2	100	1		
H. influenzae	2	50	0		
B. Friedlander	0	0	4		
B. alkaligenes	0	0	3		
B. pyocyaneus		0	. 20		
Pn. pneumoniae		0	1		

creased secretions and inflammatory edema.

From this it follows that the prevention of postoperative pulmonary infection should be directed toward:

- 1. Reduction of bronchial secretions by methods such as elimination of pre-existing infections, bronchial drainage, proper premedication, intra-tracheal anesthesia, smooth induction, and postoperative endotracheal suction.
- 2. Elimination of shallow breathing by methods such as re-breathing with carbogen mixture, frequent change of position, early ambulation, proper analysesia and suction to avoid abdominal distention.
- 3. Reduction of infection and elimination of pathogenic organisms in nose, throat and lungs by methods such as the inhalation of micronized penicillin. Micronized penicillin by inhalation seems best suited for the potentially atelectatic operative case by virtue of its concentrated local action. Numerous workers have demonstrated a progressive impairment of the circulation in a region of atelectasis, which, when coupled with the presence of infected secretions in the alveoli, would reduce the efficiency of a systemic agent such as intramuscular penicillin. Nebulized or aerosol penicillin has been disappointing in the studies of Findlay and of Holborow as well as in our small group of cases listed in Table 2. This is perhaps due to the fact that it is more rapidly absorbed from the alveoli, giving a less sustained local action than the particulate micronized penicillin. It is probable that the alveolar distribution of aerosol penicillin is less complete, due to the difficulty in controlling the inhaled droplet size, the larger droplets being deposited in the mouth and pharynx.

It should be stressed that the respiratory complications encountered in this series of cases were mild in character and did not in any instance appreciably alter the post-operative course of the patient. The one case of bronchopneumonia was in an individual with chronic basilar infection whose operation of vagotomy had been postponed previously due to this condition. He had a

temperature of 39°C. which subsided in twenty-four hours and he left the hospital on his tenth postoperative day. Four of the cases of atelectasis had fever of 38.5°C. and were turned and vigorously percussed with dislodgement of the mucous plug and a resulting fall in temperature and respiratory rate. The remaining three cases of atelectasis were entirely asymptomatic and the condition noted only on routine physical examination of the chest. No case required bronchoscopy.

Summary

- 1. A series of 150 thoracic and upper abdominal operative cases were given micronized penicillin by inhalation pre-operatively and postoperatively in an attempt to prevent postoperative pulmonary infections
- 2. There were eight complications including one case of bronchopneumonia and seven cases of atelectasis for an over-all morbidity of 5.3 per cent. This compares with a complication rate of 19 per cent in a series of untreated cases at this hospital.
- 3. At lectasis is the primary cause of postoperative pulmonary infection. Collateral ventilation prevents the development of at lectasis except in the presence of shallow breathing or pulmonary inflammation.
- 4. Shallow breathing may be eliminated by established methods.
- 5. Micronized penicillin by inhalation is an agent of considerable value in the reduction of pulmonary inflammation when used prophylactically in the operative case.
- 6. Micronized forms of some of the newer antibiotics which are active against the gram-negative organisms may further extend the usefulness of this method.

MUTUAL UNDERSTANDING

The best medical care is based on friendly, mutual understanding between physician and patient. The American Medical Association has designed an attractive new plaque for display on your office desk or wall. This plaque is an invitation to your patients to talk over questions of professional services and fees. You may secure one of these plaques for one dollar from the Order Department, American Medical Association, 535 North Dearborn Street, Chicago 10, Illinois.

CONGENITAL ATRESIA OF THE BILE DUCTS

WITH REPORT OF A CASE

PAUL D. KELLER, M.D. SALT LAKE CITY, UTAH

Reports published in the medical literature indicate that congenital atresia of the bile ducts is not common. Only 200 cases were noted in the American Medical Literature by Motsay and May when they submitted their report in 1945. At the New York Babies Hospital there were only sixteen cases in about 21,000 admissions, according to the study of Donovan. Keller and Nute analyzed the data at the St. Louis Children's Hospital and found the incidence to be only six cases in 24,441 autopsies. However, it is believed that many cases die each year, sometimes without diagnosis and often without adequate exploration.

Etiological Factors

Any explanation for the development of anomaly of the bile passages is inadequate. Certain interesting facts, however, are known. Both intra-hepatic and extra-hepatic bile ducts are formed by evagination of the foregut. During the early stages they constitute solid cords of cells. By degeneration of the central cells the lumen is eventually developed. When this process of central degeneration fails to develop, congenital anomalies result. The reports of Ladd, Holmes and of Stolkind disclose that atresia of the bile ducts may occur anywhere in the extra-hepatic portion of these structures. There are a few authentic cases, Mclendon and Graham, Parsons and Hickman, where sections of the liver show that the intra-hepatic ducts are so mal-developed that intra-hepatic biliary atresia occurs with obstruction.

Pathological Findings

According to Moore, the general variations found at autopsy or operation may be placed in the following categories: (1) patients in whom there are no demonstrable extra-hepatic ducts, (2) patients in whom there is atresia of the hepatic ducts, (3) patients in whom there is atresia of the common duct, (4) patients in whom the gallbladder is represented by a moderate-

sized cystic space not connected with the common duct, and (5) patients in whom the gallbladder connects directly with the duodenum, but in whom there are no other extra-hepatic ducts. If a child lives any appreciable length of time after birth there is invariably found an obstructive biliary cirrhosis of the liver. This organ, in a fairly advanced case, is enlarged and quite uniformly nodular. Associated acites and splenomegaly of varying degree is present.

Microscopic Pathology

There is proliferation of the intra-hepatic bile ducts. They are dilated and many of them contain accumulations of inspissated dark greenish-brown bile. There is extensive proliferation about the periportal regions of connective tissue. Phagocytic cells are usually numerous and filled with particles of concentrated bile. The usual microscopic picture is quite uniform and distinguishable from other forms of liver cirrhosis.

Clinical Features

These babies develop progressive jaundice that begins soon after they are born. The urine is dark and the stools clay-colored or white from the beginning. They usually show adequate weight-gain and their general nutritional status remains good in spite of the jaundice for several weeks. Bleeding tendencies develop late in children with congenital atresia of the bile ducts and can be avoided by giving paraenteral vitamin K.

Deep jaundice is the most prominent physical finding. A prominent abdomen with visibly dilated veins gradually develops. The liver becomes palpable and, as the case advances, the spleen also becomes palpable. Acites can be demonstrated in the older children.

The laboratory findings are those of a complete, progressive, obstructive jaundice. The most reliable test is the constant absence of urobilinogen from the urine, or its pres-

ence only in traces. The icterus index or serum bilirubin reach very high values. The Van Den Bergh reaction is of the direct type. There are no bile pigments in the stools but they are extremely concentrated in the urine. In late cases, liver function tests will show decreased function. All cases should be operated on long before there is detectable liver damage.

Differential Diagnosis

Icterus neonatorum, sometimes called physiological jaundice of the newborn, may at first be confused with congenital atresia of the bile ducts. However, this condition clears readily, beginning with the first day of birth, while the jaundice of biliary atresia steadily deepens and may not be detected until a few days after birth. Complete obstructive jaundice is present, with acholic stools, etc., in individuals with biliary atresia, while in incterus neonatorum the jaundice is of the hemolytic type.

The jaundice of erithroblastosis fetalis may, for transient periods (Davidsohn), be of the complete obstructive type. However, the presence of anemia, erthroblastemia and the usual downhill course will serve to differentiate this disease quite easily in most cases. Observations with proper study for a few days will uncover the exact nature of

Jaundice in infants with a severe bacteremia often results either from hemolysis of red blood cells or from extensive destruction of liver parenchyma. Such patients, for brief periods, may present a similar clinical picture to that of congenital atresia of the bile ducts. The former patients show the usual evidence of severe infection, viz., fever, leucocystosis, toxicity, and prostration. They are unlikely to have complete obstructive jaundice and if they do it is only for short periods.

With the modern management of syphilis and the routine practice of obtaining prenatal serology, congenital syphilis no longer offers a differential diagnostic problem.

Treatment

Early surgical exploration is the only form of treatment that offers any hope for a normal span of life to these infants. Exploration of the anomalous extra-hepatic biliary system is necessary. This requires a detailed dissection of the porta hepatis well into the hilum of the liver. Resection of a portion of the liver in search of a suitable duct for anastamosis or for the establishment of a biliary fistula should be done when no suitable remnant of the extrahepatic biliary system is present. The type of reconstructive procedure that is done will be dictated by the findings at operation. A general principle can be statedthat it is desirable to restore as nearly as possible the usual anatomical arrangement and physiological function of the normal individual.

CASE PRESENTATION

This infant was admitted to the St. Louis Children's Hospital, St. Louis, Missouri, on May 8, 1950. The history was taken from the parents and it is considered reliable:

Chief complaint: Jaundice since birth.

Present illness: The patient is an eight-weekold child that was born following a normal preg-nancy and delivery in Montgomery City, Mis-souri. Labor was thirteen hours in duration. At birth it was noticed that the child's conjunctivae were yellow and since that time there has been a progression of generalized jaundice. The father noticed that the urine is always deep yellow and he states that the stools are chalkwhite and hard. The baby has three to four bowel movements a day. With each bowel movement he screams as if he were in pain. He eats well and is on a 2:1 evaporated milk and water formula, six times a day. However, the baby frequently vomits his food ten to thirty minutes after eating. The vomitus is usually undigested food but never contains bile. The mother states that if one gently squeezes the baby's abdomen he screams as if in pain. In fact, the baby cries most of the day. Despite all this, the baby gained from five pounds and eleven ounces to eight and an half pounds since birth.

Family history: The mother and father are both 21 years old and have always been in excellent health. There is no family history of cardiac, renal, or pulmonary disease or of biliary tract disease. The mother has never received blood transfusion. They do not know their Rh

Physical Examination: The patient is an irritable, hungry, malnourished white male infant who does not appear acutely ill. Normal pulse, temperature and respiration. Skin is lemon-yellow; the conjunctivae are even deeper yellow. There are no scratch marks on the skin. No petechiae or ecchymoses are found. Mucous membranes are slightly pale despite the icterus.

Pupils react to light. The nose and throat are negative. Neck is supple with no adenopathy. Lungs, clear to percussion and auscultation. Heart tones are good and the rhythm is regular. No murmurs are heard. The abdomen is distended, with bowel pattern that is visible at times. The baby vomited twice during the draw-ing of blood. The vomitus contained no bile. The liver is two and a half finger-breadths be-low the infra-costal line. The spleen is not palpable. Extremities show no gross deformities. The right testis is twice as large as the left and somewhat firmer (hydrocele?). Reflexes are physiological. Urine: There is a deep yelloworange stain on the diaper. Stool, pale chalky and solid.

Clinical Impression: Congenital atresia of the extra-hepatic bile ducts.

Admission Laboratory Data: BBC-4,400,000 HB-8.7 grams WBC-9,400 Bleeding time—45 seconds Clotting time—2 minutes Prothrombin time-Upper limits of normal Total proteins-5.25 grams Albumen-3.88 grams Globulin-1.38 grams Cepholin flocculation-Negative Icterus index-70 units Blood culture-No growth Alkaline phosphatase-29 units RBC fragility-Normal Stool-Clay-colored-No bilirubin Urine-Bilirubin positive No urobilinogen Otherwise negative Blood type-A-Rh-Negative Kahn-Negative

Fluoroscopy of the chest was negative.

X-ray findings: AP and lateral of abdomen showed no x-ray abnormalities.



Fig. 1. Healthy child at 18 months—14 months after surgery.

Most of the laboratory studies were repeated preoperatively and found essentially as given above. Whole blood was given until the hemoglobin was within high normal limits.

On May 17, 1950, an exploratory laparotomy was done. The usual diffuse cirrhosis of the liver was encountered, with a moderate amount of acites. The spleen was enlarged about four times normal. There was no communication of the extra-hepatic bile ducts with the gastro intestinal tract. The gallbladder was a tiny cord-like structure 3 millimeters in diameter and 3½ cm. long. It joined a common hepatic duct that ended blindly with no evidence of a common bile duct.

The duodenum was mobilized and the thick-walled end of the hepatic duct stump was anastomosed to the first portion of the duodenum. A tiny polythene tube was left in the tubular gallbladder and brought out with a drain through a right upper quadrant stab wound. A liver biopsy was taken.

Gross Pathology: 50-2286 consists of three portions. No. 1 consists of specimen used for frozen section which is an irregular wedge of liver tissue approximately 1x1½x1 cm. This is greenish in color and shows fairly normal liver architecture grossly. No. 2 consists of more liver tissue, another pyramidal wedge about 2x1x1 cm. The specimen looks similar to specimen No. 1. Specimen No. 3 consists of a cyst-like structure, approximately 2x2 cm. when flattened out and less than 1 mm. thick. It is almost transparent. This is very filmy, said to be a liver cyst. Sections were made for microscopic study and labeled 1, 2, and 3, respectively. All for section. Jar O. (Bernard).

Microscopic Pathology: Sections show portal fibrosis and proliferation of bile ducts with a few ducts dilated and filled with bile. Section 3 shows a fibrous cyst wall.

Postoperative Course: The patient had an uneventful postoperative course. Light yellow bile drained from the plastic tube until it was clamped on the second postoperative day. The icterus index rapidly returned to normal and the stools took on the usual color. A cholangiogram on the sixth postoperative day showed dye to readily pass into the duodenum. The patient was discharged on the fourteenth postoperative day and has been clinically normal since.

The photograph (Fig. 1) shows the child at age 18 months, 14 months after the operation.

Discussion

Anomalies of the extra-hepatic bile passages occur often enough to warrant familiarity of the clinical picture by all who are in the practice of medicine. Although such patients need not be considered as surgical emergencies it is important to make an early diagnosis and to see that the proper management is undertaken. The indication for laporatomy with thorough exploration of the anomalous biliary system is clear in all cases. It is noteworthy that anomalies of the bile ducts are often associated with other developmental anomalies, especially, of the heart. More than one case has been reported in a single family.

Case Report

NEW DRUGS IN THE TREATMENT OF KERATITIS

JOHN A. EGAN, M.D. DENVER

For considerably more than a year, cortisone has been used topically in the eye, especially in infections of the anterior segment. Many successes have been described in several articles. So little understood are the actions of the steroids that it is impossible to evaluate the result of treatment on any selected case. Disease processes which appear similar clinically will confound us, some by responding to cortisone promptly and efficiently, and others showing absolutely no improvement.

Almost nothing has been written about the combined use of cortisone drops and antibiotic drops in the eye. From the absence of any worthwhile knowledge concerning this form of treatment, one might even wonder whether cortisone and the antibiotics are compatible. However, the following case history indicates that they

CASE HISTORY

Dr. H. M. H. was seen August 29, 1951, at about 5 p.m. with a painfully inflamed right eye of several days duration. The left eye was normal. The history given indicated that the infection had started August 25 and was progressively growing worse. He reported that he had used "a couple of drops of cortisone and one or two drops of aureomycin" during this period.

Examination revealed a deep, solid, gray-white area 3 mm. in diameter in the center of the cornea. It strained intensely with fluorescein as seen with the bio-microscope. The epithelium over this area was gone, yet the lesion showed no crater-like concavity so typical of corneal ulcer. Around the lesion, tiny and faintly staining, were pin-head punctate spots of infiltration, perhaps half a dozen in number. These were suggestive of the asterisk-like lesions noted in epidemic kerato-conjunctivitis. The larger lesion appeared to extend through the deepest layers of the cornea. Large lattice-like folds of Descemet's membrane were noted. The pupil was small and mobile. No keratic precipitates were seen, nor was there any haze in the anterior chamber. There was slight edema of the entire corneal epithelium. In addition, the eye showed all concomitants of severe keratitis such as chemosis, ciliary injection, etc. The vision was I made a tentative diagnosis of disciform keratitis.

Course: Since all bacteriological laboratories were closed, it was decided to start treatment at once without further diagnostic aid. Cortisone drops (Schering's Cortogen) were used, two in the eye every hour. At the same time two drops of aureomycin (Lederle) were also instilled. Atropine sulphate eye drops were given at once, and hot compresses advised fifteen minutes out of every hour. Between treatments the eye was patched. Chloromycetin (chloramphenicol), 250 mg., was given by mouth every four hours.

The cortisone and aureomycin drops were given five times prior to the patient's retiring. Upon arising at 5 a.m. the drops were again started and were continued hourly until I saw the patient the following morning at 10 a.m. To my amazement, on examining the eye with a strong light and using the loop, I could see no evidence of a corneal lesion. In fact, except for the pupillary dilatation, the casual observer would have pronounced the eye normal. Upon staining. With Fluorescein I detected a microscopically thin broken line of green staining in the epithelium as the only remaining evidence that the eye had been severely affected. The vision was 20/25 corrected.

Discussion

In my years of practice as an ophthalmologist, I can state that this practical recovery in approximately seventeen hours of severe keratitis was as surprising and dramatic as anything I have ever encountered. Such a condition before the advent or cortisone and antibiotics ordinarily would have taken weeks and months to recover. No one can say whether any one of the three drugs, chloromycetin, aureomycin or cortisone, was chiefly responsible for curing this patient. My inclination is to believe that all three probably helped. Ample proof exists that cortisone drops and aureomycin drops used separately are often amazingly efficient in combatting corneal infections. Recent investigative work also indicates that chloromycetin orally penetrates into the eye better than any other antibiotic. Therefore, there is no reason to believe that all three drugs were not partially responsible for this cure.

It is regrettable that we do not have sufficient cases to evaluate each of these drugs separately. Until we do, or until further information is available, I feel that we are obligated to use the most efficient and powerful drugs we have, even if done in a shotgun manner, especially when they produce such rapid and amazing cures as they have done in this case.

Organization

National Affairs

Proceedings

Programs - Society Notices - News

This Is Late, but Too Good to Miss!

Most magazines and medical journals would abhor the thought of publishing as news some-thing that happened six months ago.

But when there comes to light pictorial proof that the President of the American Medical Association was caught with his aim down—on a Wyoming antelope hunt last September—we think this bit of humor makes news even the following February.



Prexy goes a-hunting: "Wonder if this is a sufficiently dignified squat for a President of the A.M.A., but -Why!-there's one of those pronghorn things they're talking about, now - yes, guess he's in range."



"This business of using a borrowed rifle-how the h- does this bolt work? Oh, 'tisn't a bolt at all-lever action-should have had a cartridge in the chamber ahead of time -well, here goes ..."



"*! ?-!& (@!*) ?@% (** ?? @@) ()!*()... missed the !*@??..." (the remaining quotes are censored by Order of the Committee on Blood Pressure).

History records that Dr. John W. Cline of San Francisco was the honor guest at the Annual Session of the Wyoming State Medical Society Session of the wyoming State Medical Society in Rock Springs in September, 1951. Wyoming doctors arranged a post-meeting antelope hunt just north of Rock Springs, for him and other guests, including Drs. Walter Freeman of Washington, D. C., and Ralph Stuck of Denver. Later they reported a good time was had by all, but no antelope were bagged.

Now it develops that Dr. Freeman was a can-did camera fan and throughout some crucial moments of the hunt was stationed close to President Cline. Later he sent prints to Dr. Stuck, who lent them to your Editors. Neither Freeman nor Stuck vouch for the scientific accuracy of the quotes accompanying our reproductions, but

the general idea is there!

COLORADO

State Medical Society

Component Societies Elect New Officers

Many of the County and District Medical Societies in Colorado choose and install their new officers each January, although others elect or install at other times of the year. Since so many still change administrations at the beginning of each calendar year, February and the annual Directory Issue is a good time to list them. Current officers of component societies as reported to the State Society's Executive Office are:

Arapahoe County-John Simon, Jr., President; W. Hogan, Secretary.

Boulder County—Harry Jones, President; Roy Wolfe, President-elect; B. E. Peterson, Secretary

Chaffee County-C. Rex Fuller, President; E. C. Budd, Secretary.

Clear Creek County-G. E. Mason, President; L. H. Goad, Secretary.

Delta County—J. H. Humphries, President; Woodrow Brown, Vice President; R. A. Un-President; derwood, Secretary.

Denver Courty—Kenneth C. Sawyer, President; James M. Perkins, Secretary.

Eastern Colorado-H. M. Hayes, President; Jerome L. Keefe, Secretary

El Paso County—Irving H. Schwab, President; J. W. McMullen, Treasurer and President-Elect; Vernon Bolton, Vice President; E. C. Crouch, Secretary.

Fremont County—Neill B. McGrath; President; Lawrence C. Perry, Vice President; G. C. Christie, Secretary.

Garfield County—Paul J. White, President; Patience Noecker, Secretary.

Huerfano County—P. G. Mathews, President James M. Lamme, Jr., Secretary. Lake County—V. E. Kelley, President; Robert

Anderson, Secretary,



Banthine—a true anticholinergic drug with an adequate range of safety—is now made available to the medical profession in parenteral form, for use intravenously or intramuscularly in those conditions characterized by nausea and vomiting, when oral medication cannot be retained and when a prompt action is desirable.

Through its anticholinergic effects, Banthine inhibits excess vagal stimulation and controls hypermotility. In Peptic Ulcer—the value of the oral form of Banthine is now well established. However, edema in the ulcer area may indicate parenteral Banthine until the healing processes have reduced the edema.

In Pancreatitis—it has been found that parenteral Banthine relieves pain, effects a fall in blood amylase and produces a general improvement in the patient's condition.

In Visceral Spasm—it inhibits motility of the gastrointestinal and urinary tracts.

Parenteral BANTHINE is supplied in serumtype ampuls containing 50 mg, of Banthine powder. Adult dosage is generally the same as with Banthine tablets,



RESEARCH IN THE SERVICE OF MEDICINE SEARLE

Larimer County—E. Minor Morrill, President; Blair Adams, Vice President; William F. Powers, Secretary.

Las Animas County—David Barglow, President; J. E. Donnelly, Secretary.

Mesa County—Edwin R. Orr, President; R. F. Hall, Secretary.

Montrose County—T. O. Plummer, President; Ross D. Luther, Secretary.

Morgan County—V. E. Wohlauer, President; L. C. Lusby, Vice President; F. E. Roark, Secre-

Northeastern Colorado—H. P. Linton, President; Kenneth H. Beebe, Secretary.

Northwestern Colorado—F. E. Willett, President; Ligon Price, Secretary.

Otero County—John A. McDonough, President; W. R. Sisson, Secretary.

Prowers County—C. T. Knuckey, President; Edwin C. Likes, Secretary.

Pueblo County—J. H. Woodbridge, President; F. G. Tice, Jr., Secretary; Clifford F. Bramer, Treasurer.

San Juan Basin—Edward G. Merritt, President; J. G. McKinley, Secretary.

San Luis Valley—W. S. Keyting, President; C. F. Knobbe, Secretary.

Washington-Yuma Counties—C. J. Bennett, President; P. D. Keller, Secretary.

Weld County—H. E. Haymond, President; F. J. T. Roukema, Secretary.

Component Societies

NORTHEAST COLORADO

Mr. John Vance of Denver, Executive Director of Colorado Medical Service, Inc., the Blue Shield organization, was guest speaker at the January meeting of the Northeast Colorado Medical Society held January 10, 1952, at the Sterling Country Club. He explained studies now under way toward possible future development of a new service contract, under procedures outlined by the House of Delegates of the State Society last September, and answered questions from the members regarding operations of the Blue Shield plan. The next meeting of the Society will be held in Ovid on February 14.

EL PASO COUNTY

Dr. M. F. Schafer gave a comprehensive review of the functions of the various departments comprising the City-County Health Unit of El Paso County and Colorado Springs at the January 9 meeting of the El Paso County Medical Society. At this meeting applications were filed by four prospective new members, Drs. Will P. Pirkey, R. E. Meatheringham, John C. Hays, and D. Joseph Budge.

Obituaries

SOLOMON S. KAUVAR, M.D.

Dr. Solomon S. Kauvar was born in Denver, Colorado, July 30, 1910, and died December 25, 1951, after an intermittant and protracted illness due to a brain tumor. He received his medical education at the University of Chicago and was licensed to practice medicine in Colorado in 1936.

His specialty was internal medicine for which he had had excellent preparation at the University of Chicago and graduate work in New York City. He also had graduate work in Wilhelmena Hospital, Amsterdam, Holland.

He was a man of strong personality and thorough scientific training for his special field of practice. The service for which he was most widely known was as Chairman of the Denver Health and Hospital Board, appointed by Mayor Newton in 1948. His service was the initiation of the more progressive efforts along the line of medical education and public health by Denver General Hospital.

His death so early in life was considered a great loss to the practice of medicine in Denver and Colorado.

CHARLES W. KESTLE, M.D.

Dr. Kestle was born in Cripple Creek, Colorado, May 21, 1907, and was educated in the University of Colorado School of Medicine, where he received his degree of Doctor of Medicine on June 15, 1931. He received his Bachelor's Degree from the University of Denver.

In 1931 he was licensed to practice medicine in Colorado and practiced in Cripple Creek until

Dr. Kestle will be remembered as an associate of the well-known Drs. W. B. and Alexander Craig, Denver, with whom he practiced until he opened an office for himself.

In 1945 he moved to Pueblo and became a member of the Pueblo County Society. He left Colorado in 1949 and was located in Stockton, California, until the time of his death.

EMMETT VANCE GRAHAM, M.D.

Dr. Graham was born in Lee County, Virginia, on December 19, 1870, and died in Denver, Colorado, on January 17, 1952. He graduated at Emory and Henry College and the Kentucky School of Medicine and received his degree to practice in 1902. He came to Colorado Springs in the same year and moved to Denver in 1904, where he interested himself in mining activities in and about Silver Plume. He later moved to Breckenridge in 1931.

Among the things of greatest interest of his life were his activities as a country doctor in Summit County during World War I and his faithfulness to the duties as an examiner for the draft board in this sparsely settled section of the state.

In 1920 and 1921, Dr. Graham did graduate work in gynecology and obstetrics at the University of Pennsylvania, after which time he devoted himself more or less to this specialty in Denver. He was a member of the Denver County, State, and American Medical Associations.

ROBERT LEE OWENS, M.D.

Dr. Robert Lee Owens, an honorary member of the El Paso County Medical Society, died on November 22, 1951, in Lubbock, Texas. Born in Dwarf, Kentucky, on August 4, 1880, he received his medical degree in 1906 from the Hospital College of Medicine of the University of Louisville.

He had practiced in Texas and had served in World War I when he came to Colorado Springs for his health in 1920. For twenty years he



"Premarin"—a naturally occurring conjugated estrogen which has long been a choice of physicians treating the climacteric—is earning further clinical acclaim in the treatment of functional uterine bleeding.

The aim of estrogenic therapy in functional uterine bleeding is to bring about cessation of bleeding, and to produce subsequent regulation of the cycle. Once hemostasis is achieved, the maximum daily dosage of "Premarin" must be continued to prevent recurrence of bleeding. This schedule forms part of cyclic estrogen-progesterone treatment for attempted salvage of ovarian function.

"Premarin" contains estrone sulfate plus the sulfates of equilin, equilenin, β-estradiol, and β-dihydroequilenin. Other α-and β-estrogenic "diols" are also present in varying amounts as water-soluble conjugates.



An "estrogen of choice for hemostasis is 'Premarin' in tablets of 1.25 mg. ...

The usual dose for hemostasis is 2 tablets three times a day. If bleeding has not decreased definitely by the third day of treatment the dosage level may be increased by 50 per cent."*

*Fry, C. O.: J. Am. M. Women's A. 4:51 (Feb.) 1949

"PREMARIN"

Estrogenic Substances (water-soluble) also known as Conjugated Estrogens (equine)

Four potencies of "Premarin" permit flexibility of dosage: 2.5 mg., 1.25 mg., 0.625 mg., and 0.3 mg. tablets; also in liquid form, 0.625 mg. in each 4 cc. (1 teaspoonful).

Ayerst, McKenna & Harrison Limited 22 East 40th Street, New York 16, N. Y. carried on part-time practice, limiting his work to dermatology. His health finally forced his retirement in 1942 and he made his home in Texas till the time of his death.

Dr. Owens is survived by his widow and one son.

PUEBLO CLINICS

FRIDAY, APRIL 4, 1952

9:00-10:30 A.M.—Registration, Minnequa University Club.

Morning Session

Dr. J. H. Woodbridge, Presiding Chairman

10:25-Welcoming Address.-Dr. L. L. Ward.

10:30-10:45—Nasal Obstruction in Every-Day Practice.—Dr. Andrew E. Demshki, Jr.

10:45-11:05—Virus Pneumonia.—Dr. John W Gardner.

11:05-11:15—Discussion—Opened by Dr. R. D. Schilling.

11:15-11:30—Carcinoma of the Throat.—Dr. Albert McC. Tipple.

11:30-11:40—Discussion—Opened by Dr. Wm. D. Grant.

12:00-2:00-Recess.

Afternoon Session

Dr. Frederick G. Tice, Jr., Presiding Chairman

2:00-2:20—Total Hysterectomy for Benign Uterine Condition.—Dr. Scott A. Gale.

2:20-2:30—Discussion—Opened by Dr. Jesse W. White.

2:30-2:50—Baby's First Year.—Dr. John Yeager. 2:50-3:00—Discussion—Opened by Dr. Ray Taylor, Jr.

3:00-3:30-Recess to view exhibits.

3:30-3:50—Hernia—Newer Aspects of Repair.— Dr. George P. Cribari.

3:50-4:00—Discussion—Opened by Dr. Eugene B. Ley.

4:00-4:20—Bladder Complications and Care Following Surgical Procedures in General.—Dr. William C. Shontz.

4:20-4:30—Discussion—Opened by Dr. George M. Meyers.

4:30-5:30-Review of exhibits.

Evening Session

6:30-Preview, Lounge.

7:00-Dinner, Dining Room.

"Office Gynecology"—Dr. Walter J. Reich, Guest Speaker.

WYOMING State Medical Society

News Notes

The Wyoming State Medical Society will hold its Annual Session in Lander on June 5, 6, and 7, 1952. The scientific meetings and exhibitors will be at the State Armory Building in Lander and the hotel headquarters will be the Noble Hotel in Lander.

COLORADO

State Health Department

MATERNAL GESTATION CALCULATORS AVAILABLE

Recently the Maternal and Child Health Section of the Colorado State Department of Public Health has received a limited number of special cardboard "slide-rules" from the U. S. Public Health Service and Children's Bureau. These are being distributed to all Colorado hospitals to assist physicians in the accurate reporting of length of pregnancy in weeks on the birth certificate.

Although most authorities agree that the birth weight is the most reliable single index of maturity, the recent emphasis on accurate recording of birth weight on all birth certificates should not be interpreted as implying that information on the length of gestation is no longer of importance. The latter is extremely valuable to physicians and health departments concerned with the reduction of infant mortality, immunity (prematurity) and related problems of the newborn.

A recent pamphlet* issued by the National Office of Vital Statistics makes the following comments regarding length of pregnancy: "For statistical purposes, period of gestation is defined as "the number of completed weeks that have elapsed between the first day of the last menstrual period and the date of birth of the child.' It is recognized that there are cases where the exact date of the mother's last menstrual period cannot be determined accurately. However, the bias introduced by such errors will be relatively minor. From a statistical standpoint, the important factor is the consistent application of the definition on the basis of the best available evidence.

"At the present time, important innaccuracies are evident in gestational information derived from birth records. These arise principally from difficulties in computing length of pregnancy in weeks, and are reflected in the statistics by extreme heaping at thirty-six and forty weeks and a minor bias toward other even weeks. The heaping at thirty-six weeks is the result of incorrectly converting nine calendar months (full-term gestation age) into weeks by considering four weeks the equivalent of a month. Heavy concentrations at forty weeks are indicative in part of a failure to calculate period of gestation for the newborn infants who seem to be normally developed."

All Colorado physicians are urged to give this matter their thoughtful attention. The gestational calculators are offered as a means of simplifying and standardizing the procedure.

*Recommendations for Developing Comparable Statistics on Prematurely Born Infants and Neonatal Mortality—National Office of Vital Statistics and the U. S. Public Health Service, Children's Bureau.

The technics used in tuberculosis control among the Indians parallel those among non-Indians and are flexible enough to be changed as accepted methods are improved. However, additional methods of attack are used that are not now considered necessary in the general population, such as the wide-scale use of BCG vaccine.—H. DeLien, M.D., and Arthur W. Dahlstrom, M.D., Am. J. Pub. Health, May, 1951.

from among all antibiotics, Neurologists and Neurosurgeons often choose

AUREOMYCIN

because


It readily passes into the cerebrospinal fluid, the presence of meningitis making little difference in its concentration.

Measurable serum levels are maintained for as long as 12 hours after oral administration, oral doses of 5 to 10 mg. per kilo at 6-hour intervals being adequate for this purpose. Aureomycin has been shown to be highly effective against those bacterial invaders commonly encountered in central nervous system infections.

Aureomycin has been reported to be effective against susceptible organisms in: Brain Abscess • Cranial Trauma Infection • Encephalitis • Meningitis

Throughout the world, as in the United States, aureomycin is recognized as a broad spectrum antibiotic of established effectiveness.

Capsules: 50 mg.—Bottles of 25 and 100; 250 mg.—Bottles of 16 and 100. Ophthalmic: Vials of 25 mg. with dropper; solution prepared by adding 5 cc. of distilled water.



LEDERLE LABORATORIES DIVISION

AMERICAN Granuld courses.

30 Rockefeller Plaza, New York 20, N.Y.

COLORADO

Medical School Notes

Dr. Ward Darley, Director of the University of Colorado Medical Center, has been appointed to the National Advisory Mental Health Council, a council of twelve members which reviews all applications for research grants in its field from the U. S. Public Health Service. Announcement was made January 22 by Surgeon General Leonard A. Scheele in Washington.

The University of Colorado School of Medicine is pioneering a new trail in postgraduate medical education. It is sending twelve members of its faculty to Cheyenne, Wyoming, where they will conduct a series of twelve lectures designed to bring family doctors in Wyoming up-to-date on the latest medical advances.

The course is the first ever conducted by the school outside of Colorado. It is being co-sponsored by the C. U. Office of Graduate and Postgraduate Medical Education and the Wyoming Chapter of the American Academy of General Practice. Lectures will be held weekly from 7:30 p.m. to 9:30 p.m. every Wednesday in the Veterans' Administration Hospital in Cheyenne. The course consists of: February 20—"Diagnosis and Management of Diabetes and Diabetic Coma," by Dr. Robert W. Gordon; "Frequent Errors in Gynecological Surgery," by Dr. E. Stewart Taylor. February 27—"Diagnosis of Common Chest Conditions, Plus Recognition of Some of the Less Common Conditions," by Dr. Robert S. Liggett;

"Treatment of Fractures by General Practitioners," by Dr. John T. Jacobs. March 5—"Management of Ulcerations of the Gastro-Intestinal Tract," by Dr. Frank B. McGlone; "Initial and Follow-Up Treatment of Severe Burns," by Dr. Mordant E. Peck. March 12—"Recognition of Complicating Factors in Pregnancy," by Dr. Freeman H. Longwell; "Diagnosis of Common Neurological Conditions," by Dr. G. Milton Shy. March 19—"Treatment of Hernia, Inguinal, Umbilical and Post-Operative, With Discussion of the Use of the Silver Wire Screen," by Dr. Edgar W. Barber; "Fluid Balance in the Very Ill Patient, Surgical and Medical," by Dr. MacDonald Wood. March 26—"Surgical Treatment of Vascular Lesions of the Lower Extremities," by Dr. Henry Swan II; "Anesthesia, Use of Sodium Pentothal, Curare, N20 and 02, Maintaining Blood Pressure at Normal Levels," by Dr. Robert W. Virtue.

The course actually is a practical review for general practitioners. The instructors, in addition to being faculty members, all are in active practice. The courses will be approved for twelve hours of credit by the Wyoming Chapter of General Practice.

WINTER AND SPRING POSTGRADUATE COURSES, 1952

Make Your Plans in Advance

Internal Medicine for General Practitioners; March 20, 21, 22, 1952—This is a practical course devoted to recent advances in diagnosis and treatment of common medical diseases. Emphasis will be placed on individual bedside and conference case discussions of selected patients from the wards in Colorado General and Denver Gen-

We Believe --

That Professional Men should be consulted on problems of sickness and health.

That Professional Men should be consulted on problems of investments.

CONSULT YOUR INVESTMENT BANKER

MAin 6281

Peters, Writer & Christensen, Inc.

724 Seventeenth Street

Denver 2, Colo.

211 Association Bldg.

Loveland, Colo.


Phone Loveland 302

Investment Bankers



All Children Can Benefit from this Protective Hot Drink at Breakfast

In its widely distributed leaflet No. 268, "Eat a Good Breakfast," the U. S. Dept. of Agriculture states: "Summer or winter, there's something hot, as a rule, in a good breakfast.... Something hot is cheering and tones up the whole digestive route."



A cup of hot Ovaltine makes an excellent contribution of virtually all essential nutrients, adding substantially to the nutritional start for the day. It also serves in a gustatory capacity by enhancing the appeal of breakfast and making other foods more inviting.

The nutrient contribution made by a cup of Ovaltine is apparent from the table below. Note the wealth of essentials added to the nutritional intake by making the simple recommendation of adding a cup of hot Ovaltine to the child's breakfast.


THE WANDER COMPANY, 360 N. MICHIGAN AVE., CHICAGO 1, ILLINOIS



Ovaltine

Here are the nutrients that a cupful of hot Ovaltine, made of ½ oz. of Ovaltine and 8 fl. oz. of whole milk,*provides:

			b. a man.
PROTEIN	10.5 Gm. IRON	4 mg.	NIACIN 2.3 mg.
			VITAMIN C 10 mg.
			VITAMIN D 140 I.U.
			CALORIES 225
PHOSPHORUS	315 mg. RIBOFLAVI	N 0.7 mg.	*Based on average reported values for milk,



in lobar pneumonia:

The prompt response to Terramycin therapy in lobar pneumonia is consistent with results obtained in primary atypical pneumonia, bronchopneumonia and many other infections of the respiratory tract. In a typical series of pediatric cases, Terramycin-treated, "temperatures returned to normal in 24 to 48 hours after therapy was begun. The clinical appearance of marked improvement took place during the same period."

Potterfield, T. G., and Starkweather, G. A.: J. Philadelphia General Hosp. 2:6 (Jan.) 1981

ANTIBIOTIC DIVISION



Terramycin is also indicated in a wide range of

GRAM-POSITIVE BACTERIAL INFECTIONS

Lobar pneumonia · Mixed bacterial pneumonias

Bacteremia and septicemia

Acute follicular tonsillitis

Septic sore throat . Pharyngitis

Acute and chronic otitis media

Acute bronchitis . Laryngotracheitis

Tracheobronchitis . Sinusitis

Chronic bronchiectasis

Pulmonary infections associated

with pancreatic insufficiency

Scarlet fever . Urinary tract infections

Acute and subacute purulent conjunctivitis

Acute catarrhal conjunctivitis

Chronic blepharoconjunctivitis

not involving the meibomian gland

Abscesses · Cellulitis

Furunculosis • Impetigo

Infections secondary to Acne vulgaris

Erysipelas · Peritonitis

GRAM-NEGATIVE BACTERIAL INFECTIONS

Gonorrhea . Brucellosis

Bacteremia and septicemia

Friedländer's pneumonia

Mixed bacterial pneumonias

Pertussis • Diffuse bronchopneumonia

Post-partum endometritis . Granuloma inguinale

Dysentery • Urinary tract infections

Respiratory tract infections

Cellulitis · Peritonitis · Tularemia

SPIROCHETAL INFECTIONS

Syphilis . Yaws . Vincent's infection

RICKETTSIAL INFECTIONS

Epidemic typhus . Murine typhus

Scrub typhus . Rickettsialpox

Q fever · Rocky Mountain spotted fever

VIRAL INFECTIONS

Primary atypical pneumonia (virus pneumonia)

Lymphogranuloma venereum · Trachoma

PROTOZOAL INFECTIONS

Amebiasis

Available as

CAPSULES

ELIXIR ORAL DROPS

INTRAVENOUS

INTRAVEROUS

OPHTHALMIC

OIMIMENT

OPHTHALMIC SOLUTION

302011014

took

: 952

tent

cal

act.

S

many

_

CHAS. PFIZER & CO., INC., Brooklyn 6, N. Y.



Jiggs Gets Four "Hot-Foots"

Cappy Miller's bought himself a new car. We won't be seeing his old jalopy bouncing over the back roads any more. I'm going to miss it, too.

Many a morning Cappy and I drove off in that rattletrap for a day's hunting or fishing. We'd pile rods or guns in back, and prop open the trunk compartment—so Jiggs, Cappy's pointer, could jump in and go along.

They say when Cappy brought the new car home he opened up the hood to show off the engine—and poor old Jiggs hopped right in! Figured it was the trunk. He hopped right out in a hurry, too. That cylinder head was mighty hot.

From where I sit, old habits are hard to shake, once they get a hold. For instance, too many people are still in the habit of trying to run their neighbors' lives—telling them how to act, how and where to practice their profession, whether or not to enjoy a glass of beer. I say that kind of thinking's outmoded...ought to be turned in for a new model!

Joe Marsh

Copyright, 1952, United States Brewers Foundation

eral Hospitals. Professor Gordon Meiklejohn and Staff. Enrollment limited to twenty-five students.

Gynecology, Obstetrics, and Related Problems of the Newborn: April 10, 11, 12, 1952—This three-day refresher course will be high-lighted by four prominent guest lecturers, Prof. Emil G. Holmstrom, University of Utah; Prof. William C. Keettel, University of Iowa; Prof. Gilbert J. Vosburgh, Western Reserve University, and one guest pediatrician. These men, together with the faculty of the University of Colorado, form a teaching team of outstanding authorities. Time will be set aside for general discussion and questions where there can be a free exchange of ideas by the participants in the course.

Poliomyelitis: May 1, 2, 3, 1952—This course is planned to review the diagnosis and management of patients with poliomyelitis. This disease presents an increasing medical problem and every physician is called upon to answer questions and advise patients and their families concerning this disease. The instructors in this course have had wide experience in managing poliomyelitis cases during the recent outbreak in Colorado.

Traumatic and Emergency Surgery: May 19 and 20, 1952—This course is for both general practitioners and specialists and will include discussions in fractures, burns, shock, antibiotics, and other drugs, transfusions, and acute abdominal conditions. This course is designed to review recent developments of these various subjects by lectures, demonstrations and non-operative clinics. The basic anatomical and physiological principles underlying these subjects will be emphasized.

Psychiatry for General Practitioners: June 26, 27, 28, 1952—This course aims to present prevalent psychiatric concomitants in general medicine. Emphasis will be placed on psychotherapy and adequate time will be provided for discussion. Guest lecturer will be Dr. William T. Shanahan, Professor of Psychiatry, University of Texas, Galveston.

Applied Medical Science Courses: January 2 through June 9, 1952—This is a regular full-time course designed to orient the graduate student in the basic sciences required for certification in the various American Specialty Boards. The following part-time courses may be arranged: Surgical Pathology, two one-half days weekly from March 24 through June 9. This will be a systematic review of surgical pathology with emphasis on the more common lesions and each system will be covered as completely as the time allows. Surgical Anatomy, four one-half days weekly from March 24 through June 9. This course is planned to be of special interest to physicians who are preparing for speciality boards. Electrocardiography, a two-hour lecture, 7:30 to 9:30 each Thursday evening beginning January 17 through March 13. Dr. Abe Ravin. Biophysics, every Monday, beginning January 14 through February 18, 7:00 to 9:00 p.m. Industrial Medicine, every Tuesday, beginning January 22 through March 11, 9:00 to 10:30 p.m.

Are you keeping up with rapid advances in medicine? These courses are designed to help you. For further details and registration write to Postgraduate Education, University of Colorado School of Medicine, 4200 East Ninth Avenue, Denver, Colorado. The above courses will be held at the University of Colorado Medical Center, Denver.

for February, 1952

141

EARNEST DRUG COMPANY

T. H. BRAYDEN, Prop.

PRESCRIPTION SPECIALISTS

1699 Broadway

Phone KEystone 7237

Denver, Colorado

"Conveniently Located for the Doctor"

and

EARNEST DRUG DISPENSARY

(Successors to Carey Drug Dispensary)

Located in the Majestic Building

217 16th Street

Phone KEystone 3265

Prompt Free Delivery Service From Both Stores

From 10 A.M. to 8 P.M.

The BROWN SCHOOLS

For Exceptional Children

Four distinct units. Tiny Tots through the Teens. Ranch for older boys. Special attention given to educational and emotional difficulties. Speech, Music, Arts and Crafts. Full time Psychologist. Under the daily supervision of a Certified Psychiatrist. Registered Nurses. Private swimming pool, fireproof building. View Book. Summer Camp. Approved by State Division of Special Education.

BERT P. BROWN

President

Paul L. White, M.D., F.A.P.A., Medical Director

P. O. Box 4008, Austin, Texas

NEW MEXICO

Medical Society

Obituaries

IRA L. PEAVY, M.D.

Dr. Ira L. Peavy, Santa Fe, died December 25, 1951, at his home following a lingering illness. At the time of his death Dr. Peavy was Director of the Venereal Disease Division of the State Department of Public Health.

Dr. Peavy was born June 14, 1889, and graduated from the University of Colorado in 1924.

A veteran Colorado and New Mexico physician, Dr. Peavy had been with the Health Department some ten years, with time out between 1945 and 1947 for work with the Colorado Fuel and Iron Company at Valdez, Colorado. Before that time he was with the St. Louis Rocky Mountain Fuel Company in Raton. He joined the State Health Department in 1942 as V-D clinician in District 6 at Carlsbad.

After returning from Colorado in May, 1947, he assumed a similar post in Las Vegas and later in Santa Fe, he became head of the division on February 1, 1951. He was a member of San Miguel County Medical Society, the New Mexico Medical Society, and the American Medical Association.

C. LEROY BROCK, M.D.

Dr. C. Leroy Brock, Albuquerque, died January 5, 1952, after a long illness. Dr. Brock had practiced in Albuquerque for thirty years, and practiced in Espanola prior to that time.

He was born in 1884 and graduated from Georgetown University in 1911. He was a member of Hope Lodge No. 20, A. F. & A. M., of Washington, D. C., the Scottish Rite Bodies at Santa Fe, and Ballut Abyad Temple Shrine of Albuquerque, and of the Episcopal Church. He was a member of Bernalillo County Medical Society, the New Mexico Medical Society, to which he was elected an Emeritus Member in 1950, and of the American Medical Association.

BLUE CROSS and BLUE SHIELD

RESULTS OF SENATE SURVEY

A Senate Subcommittee, under the chairmanship of Herbert H. Lehman, former Governor of New York, has recently completed a study of Health Insurance Plans in the United States. A summary of the report is published by the Spectator, Property Insurance Review', for September, 1951, and here we present a summary of this summary. Some of the figures go back a year or two, since the study has been under way for considerable time.

The number of people carrying health insurance in the United States is 75 million—about one-half of the population. Blue Cross protects 37 million, Blue Shield 18 million, insurance companies 34 million. There is some duplication



A clinical record stretching over 17 years of use in a wide variety of procedures . . .

1803 reports published in medical journals throughout the civilized world . . . this is the background Pentothal Sodium offers today's anesthesiologist. Through the years, the Abbott-discovered ultra-short-acting barbiturate has become virtually synonymous with intravenous anesthesia. For good reasons. There is a rapid, pleasant induction, complete surgical amnesia. The patient usually awakens without nausea. With Pentothal, the explosion hazard is eliminated, the equipment simple and easily stored.

When individual requirement warrants, Pentothal may be combined with any number of other anesthetics. Investigate the full potentialities of Pentothal in minor and major surgery—and in obstetrics—by writing Abbott Laboratories,

North Chicago, Illinois, for detailed literature.

As an adjunct to PENTOTHAL Sodium

TUBOCURARINE Chloride, Abbott

... supplied in 10-cc, and 20-cc, vials, each cc. representing 3 mg. of tubocurarine chloride pentahydrate. Also 1-cc. ampoules, 15 mg. Ask for literature. Pentothal Sodium

FOR INTRAVENOUS ANESTHESIA

Cook County Graduate School of Medicine

ANNOUNCES CONTINUOUS COURSES

ANNOUNCES CONTINUOUS COURSES
SURGERY—Intensive Course in Surgical Technic, Two
Weeks, starting February 4, February 18, March 3.
Surgical Technic, Surgical Anatomy and Clinical
Surgery, Four Weeks, starting March 3, June 2. Surgical Anatomy and Clinical Surgery, Two Weeks,
starting March 17, June 16. Surgery, Or Colon and
Rectum, One Week, starting March 3, April 7. Gallbladder Surgery, Ten Hours, starting April 21. Basic
Principles in General Surgery, Two Weeks, starting
March 31. Breast and Thyroid Surgery, One Week,
starting June 23. Esophageal Surgery, One Week,
starting June 23. Thoracic Surgery, One Week,
starting June 23. Facebragery, One Week,
starting June 23. Facebragery, One Week,
starting June 23. Facebragery, One Week,
starting February 4. Weeks, starting February 4.

GYNECOLOGY—Intensive Course, Two Weeks, starting February 18, March 17. Vaginal Approach to Pelvic Surgery, One Week, starting March 3, March 31.

OBSTETRICS—Intensive Course, Two Weeks, starting March 3, March 31.

EDICINE—Intensive General Course, Two Weeks, starting May 5. Electrocardiography and Heart Disease, Two Weeks, starting March 17. Gastroenterology, Two Weeks, starting May 19. Hematology, One Week, starting June 16.

UROLOGY—Intensive Course, Two Weeks, starting April 28. Ten-Day Practical Course in Cystoscopy, start-ing February 18, March 3, and every two weeks.

ROENTGENOLOGY—Two-Week Lecture and Clinical Courses each month.

GENERAL, INTENSIVE AND SPECIAL COURSES IN ALL BRANCHES OF MEDICINE, SURGERY AND THE SPECIALTIES

TEACHING FACULTY—ATTENDING STAFF OF COOK COUNTY HOSPITAL ADDRESS: REGISTRAR, 427 SOUTH HONORE STREET, CHICAGO 12, ILLINOIS

ACCIDENT - HOSPITAL - SICKNESS INSURANCE

Physicians, Surgeons, Dentists Exclusively



\$15,000.00 accidental death \$20,000.00 accidental death \$32.00 \$100.00 weekly indemnity, accidet and siekness
Cost has never exceeded amounts shown. ALSO HOSPITAL POLICIES FOR MEMBERS, WIVES AND CHILDREN AT SMALL ADDITIONAL COST.

85c out of each \$1.00 gross income used for members' benefit

\$4,000,000.00 \$17,000,000.00 INVESTED ASSETS PAID FOR CLAIMS

220,000.00 deposits with State of Nebraska for protection of our member.
Disability need not be incurred in line of duty—
benefits from the beginning day of disability.
PHYSICIANS CASUALTY ASSOCIATION
PHYSICIANS HEALTH ASSOCIATION

49 years under the same management 100 First National Bank Building, Omaha 2, Nebraska

since 31 million people carry both hospital and surgical insurance. Between three and four million carry relatively complete insurance. Blue Cross pays 70-80 per cent of the hospital bill for which they are responsible; the insurance companies pay 45-55 per cent. Blue Shield pays 67-75 per cent of surgical and medical expense; the insurance companies pay 46 per cent.

Subscription payments to Blue Cross in 1949 amounted to 303 million dollars; to Blue Shield 79 million; to insurance companies and independent plans 373 million. These are merely insurance payments; the total spent in 1949 for health services exceeded 10 billion dollars. According to the Senate report employers pay about one-half of the premiums in insurance plans under collective bargaining agreements.

Government funds provide 71 per cent of hospital beds, many of these beds taking care of chronically ill patients. State governments provide 45 per cent of these beds, the federal government 13 per cent, local governments likewise 13 per cent. One-fourth of patients admitted to hospitals enter government institutions.

Activities of the federal government include free medical care to veterans, grants to states for infant care, community inocculations, water and food inspection, free milk for grade schools, etc. State activities include the operation of hospitals, maintenance of public health departments, sanitation, venereal disease control, industrial safety, etc. Local governments usually make contact with patients and they operate hospitals and employ physicians, dentists, and nurses. In these activities they receive financial support from state and federal agencies.

Over 4.5 million persons are on the rolls of public relief agencies, the break-down running as follows: Aged, 2,625,594; dependent children, 1,365,813; blind, 89,301; general relief, 461,000. The cost of this relief in 1949 was 2,234 million

According to the American Medical Association the average cost of illness requiring hospitaliza-tion is \$285. Of this sum \$131 is allocated to the hospital; \$109 to the physician; \$10 to special nursing care; and \$14 to incidental items.

This minimal summary will show the general nature of the Senate Subcommittee's report. It appears that the study was made for purposes of information rather than propaganda but it may be anticipated that the findings of the report will yield to diverse interpretations.

³The Spectator, Chestnut and 56th Streets, Philadelphia, Pennsylvania.

²Currently 43 million.

³Currently 21 million.

The Book Corner

Book Reviews

Syllabus of Human Neoplasms: By R. M. Mülligan, M.D., Professor of Pathology in the University of Colorado School of Medicine: with 230 illustra-tions; Lea & Febiger, Philadelphia, 1951. Price,

The authoritative impression which the reader gains from the preface of this book maintains throughout. No verbosity is encountered; descriptions are terse and exact.

TO ALL MEMBERS AND GUESTS

Of the Medical Profession, Who Attend the MIDWINTER POSTGRADUCATE CLINICS

TECHNICAL EQUIPMENT CORP .-

The House Service Is Building—
Extends to You a Hearty Welcome

You are cordially invited to Booth 23—pause for a few moments—return to your lecture rooms, refreshed and ready for more of the interesting clinics you have come to see and hear.

This year our exhibit is planned solely for your entertainment. We offer no items for sale.

Of course, if you develop a deep desire for a

KELLY-KOETT X-RAY

We will accept your orders.

Our Service is COURTEOUS — FRIENDLY — PROMPT

TECHNICAL EQUIPMENT CORPORATION

2548 W. 29th Ave.

GLendale 4768

DENVER 11, COLORADO

The eleven chapters are divided in simple, logical fashion with general ideas and definitions making up the first. The latest concepts regarding epithelial neoplasms of skin and appendages are found in Chapter II. Chapter III treats of neoplasms of mesenchymal origin. The fourth chapter is very well done and has to do with neoplasms of neuroectodermal origin, and removes the old veil of uncertainty from discussion of these tumors. Chapters V, VI, VII, VIII, IX, and X are concerned with neoplasms of the alimentary tract, respiratory tract, urinary tract, male genitalia, female genitalia, and endocrine glands. Unusual neoplasms are described in Chapter XI and in the Supplement.

The bibliography at the end of each chapter is made up of references to recent literature for the most part and is gratifyingly complete.

Illustrations are numerous, appear characteristic, and still different from the ones usually found in pathology text books. Many of these illustrations are excellent.

Pathologists will welcome this up-to-date book to their reference libraries. It will prove valuable to surgeons and to all physicians interested in tumors.

FRANCES McCONNELL, M.D.

A Textbook on Medicine: Edited by Russell L. Cecil, M.D., Sc.D., Professor of Clinical Medicine Emeritus, Cornell University, New York; Robert F. Loeb, M.D., Bard Professor of Medicine, Columbia University, New York; Associate Editors, Alexander B. Gutman, M.D., Professor of Medicine, Columbia University, New York; Walsh McDermott, M.D., Associate Professor of Medicine, Cornell University, New York; Harold G. Wolff, M.D., Associate

A new edition of Cecil's Medicine is always an event of interest both to the internist and to the general practitioner. This book has been the standard text of the present generation of practitioners and medical students, just as Osler was the constant guide of our fathers in medicine. The new edition does not suffer by comparison with previous ones.

Dr. Cecil has had associated with him as editors, Robert F. Loeb, Alexander B. Gutman, Walsh McDermott, and Harold G. Wolff. The editors have added twenty new subjects, and yet have managed to shorten the volume by 136 pages with no sacrifice of important material.

This has reduced the thickness of the book by one centimeter, making it somewhat more easily handled. Some eighty-two new articles on subjects previously covered have been prepared.

Certain trends may be worthy of mention. There is a new section on collagen diseases. ACTH and cortisone are discussed fairly well in several sections of the book. Infectious mononucleosis is now classed as a viral disease. There is a new section on inborn errors of metabolism. Richards has an interesting discussion of pulmonary function in health and disease. Many revisions have been made in the field of endocrinology, and a recurrent theme throughout the entire volume is the importance of endocrine and emotional factors in many disease processes.

One word of criticism is in order. Colorado

One word of criticism is in order. Colorado Tick Fever is a proven and well-defined clinical entity of some years standing. It is mentioned nowhere in this book.

THEODORE K. GLEICHMAN, M.D.

Primer on Fractures: Prepared by the Special Exhibit Committee on Fractures in Cooperation with the Committee on Scientific Exhibit of the American Medical Association; Sixth Edition. Paul B. Hoeber, Inc., Medical Book Department of Harper & Brothers, 1951. Price, \$2.00.

This small (109 pages) book presents the essential principles of emergency treatment of fractures and application of plaster of paris bandages. For each of the common fractures of the body, the treatment is outlined as to reduction, immobilization and after care, with demonstration by drawings of the most accepted methods. It is regrettable, however, that the Committee on Fractures still advocates and even diagrams the use of the "banjo splint" for phalangeal fractures.

An unusual, but most practical, aspect of the primer is its blank pages on the left half of the book as space for the notes by the student or practitioner.

As it is concise and fairly complete as to the management of fractures, a student who would learn all that is contained in this short book would have little difficulty in treating almost any type of fracture.

MacDONALD WOOD, M.D.

DAVIS BROS. DRUG CO.

WHOLESALE DRUGS

1628 15th Street, Denver, Colorado

Phone KEystone 5131

MERCY HOSPITAL

Conducted by Sisters of Mercy School of Nursing in Connection

A General Hospital Scientifically Equipped

1619 Milwaukee St., Denver

FRemont 2771

DENVER TOWEL SUPPLY COMPANY

1730 Speer Blvd.

TAbor 3276

Denver, Colorado

FOR INFANT FEEDING

Baker's Modified Milk

PROVIDES A NUTRITIONALLY ADEQUATE* FORMULA

PROTEIN

Baker's Modified Milk supplies 3.5 to 4 grams of milk protein per kilogram of body weight per day when the total feeding is 2½ fluid ounces of normal strength dilution** per pound of body weight. The milk protein in Baker's also provides adequate amounts of all the essential amino acids.

CARBOHYDRATE

The carbohydrates in Baker's Modified Milk are lactose and dextrose. The dextrose which requires no digestion is readily assimilated. The lactose is slowly digested and absorbed. This combination of sugars is less likely to lead to digestive disturbances than if a single sugar were used. The carbohydrate content (7% at normal dilution**) provides adequate calories to spare the protein for its normal function of tissue building and repair.

FAT

The fat-carbohydrate ratio (approximately 1:2) is adequate to

Made from Grade A Milk (U.S. Public Health Service Milk Code) which has been modified by replacement of the milk fat with vegetable and animal fats by the addition of carbohydrates, vitamins and iron.

insure proper fat metabolism. The butter fat has been replaced by other fats containing less of the undesirable very low and very high molecular weight fatty acids. The added fats have also been selected to provide adequate amounts of the essential unsaturated fatty acids.

MINERALS

Baker's Modified Milk contains an adequate mineral content with the calcium-phosphorus ratio falling within the optimum range (1.3 to 1). Since cows milk contains only a trace of iron, sufficient iron ammonium citrate has been added to supply 7.5 milligrams of iron per quart of normal dilution.**

VITAMINS

Baker's Modified Milk supplies adequate amounts of all recommended vitamins except Vitamin C.

> These are the reasons why infants thrive on Baker's Modified Milk—a nutritionally adequate* formula.



When fed in normal quantities, provides amounts of proteins, vitamins (except C), minerals and essential unsaturated fatty acids equal to or exceeding the daily recommended allowances of the Food and Nutrition Board of the National

Research Council.
**Dilute with equal parts of water.

BAKER'S MODIFIED MILK

THE BAKER LABORATORIES INC.

Main Office: Cleveland, Ohio Plant: East Troy, Wisconsin Division Offices: Atlanta, Dallas, Denver, Greensboro, N. C., Los Angeles, San Francisco, Seattle



Clinical Heart Disease: By Samuel A. Levine, M.D., F.A.C.P., Clinical Professor of Medicine, Harvard Medical School; Physician, the Peter Bent Brigham Hospital, Boston; Consultant Cardiologist, Newton-Wellesley Hospital; Physician, New England Baptist Hospital, Fourth Edition, illustrated. W. B. Saunders Company, Philadelphia and London, 1951.

To the long list of texts of cardiology, Dr. Levine has added his fourth edition of the clinical approach to heart disease. For those who are familiar with Levine's informal style of writing, this new edition will come as no great surprise except to bring them up to date; for those others who have not yet had the good fortune to read his previous books a treat is in store. From a long and rich experience, Levine brings a form of judgment and maturity so necessary in these harried days of increasing heart consciousness. It is as if one were attending his rounds and listening to his little stories of other cases which have fooled him that so enriches his writing.

This fourth edition comes to us at an important period in the field of cardiology. It brings up to date the therapy of subacute bacterial endocardititis and includes a section on anticoagulant prophylaxis. The section on congenital heart disease has been enriched by studies with

COLVIN MEDICAL BOOKS

Medical Publications of All Publishers

Books Sent for Examination on Request

We Maintain This Book Store for

Your Convenience

Write or come to
705-706 MAJESTIC BUILDING
Denver 2, Colo. Call MAin 3866

cardiac catheterization, an understanding of which is essential for proper selection for surgery. The previously large section on electrocardiography has been further expanded to include the latest in precordial and unipolar leads as they apply in the correct evaluation of coronary disease.

As a single text on heart disease, Levine's fourth edition is felt to be one of the most rounded of all presently on the market and is heartily recommended to all interested in the field

ALLAN HURST, M.D.

Genetics in Ophthalmology: By Arnold Sorsby, Research Professor in Ophthalmology, Royal College of Surgeons and Royal Eye Hospital; Surgeon, Royal Eye Hospital; Surgeon, Royal Eye Hospital, London. Butterworth & Co. (Publishers), Ltd., London, England. The C. V. Mosby Company, St. Louis, Mo., U. S. A., 1951. Price, \$9.50.

This new work by Arnold Sorsby is one showing an endless amount of research in the field of genetics in ophthalmology. A detailed review in limited space is not possible. The text requires close study and covers the field thoroughly.

The first section deals with theoretical concepts of modes of inheritance, some general concepts, human pedigrees, clinical varieties of genetic disease, and prospects in the control of genetic disease.

The second section deals with the globe as a whole, the cornea, lens, uveal tract, optic nerve and other tissues. Section three considers metabolic disorders, some systemic disorders and syndromes. A carefully prepared bibliography is appended.

Ophthalmologists and geneticists who take the time to study this new text carefully will find it most interesting and instructive.

JAMES M. SHIELDS, M.D.

H. C. STAPLETON DRUG COMPANY

Service Wholesalers for the Prescription Department

RAPID-INTELLIGENT-SERVICE

1252-54 Arapahoe St., Denver, Colo.

Phone MAin 4152

SHIRLEY-SAVOY HOTEL

At Your Service

New Lincoln Auditorium and Private Dining Room

J: Edgar Smith, President Ed C. Bennett, Manager BROADWAY and EAST 17th AVENUE, DENVER, COLO.

Ike Walton, Managing Director TAbor 2151

St. Anthony Hospital

Write or Phone Registrar for Information

West 16th Ave. and Quitman, Denver, Colorado

AComa 1761

Cooperating With the Ethical Medical Profession

THE COLORADO ARTIFICIAL LIMB COMPANY, Inc.

Authorized Manufacturers of the Famous Rowley Legs

1437 17th Street

MAin 2866

Denver, Colo.

a new year

Brings You This New Service

The world famous Stacey's, medical book firm of San Francisco, has joined hands with the skilled organization of the Technical Book Company of Denver to form a new company — Stacey-Technical Book Company of Denver. This new working arrangement has been made in order to develop an outstanding center of medical books from which the physicians of Colorado may receive superior service. You are cordially invited to drop in and browse, or phone, or write for any of your book requirements.

Stacey - TECHNICAL BOOK CO.

1814 STOUT STREET
DENVER 1, COLORADO

A Western Institution

eeded

The DALLAS SOUTHERN **CLINICAL SOCIETY**

Announces Its

21st ANNUAL

Spring Clinical Conference

MARCH 17, 18, 19, 20, 1952

for information address:

Executive Secretary 422 Medical Arts Bldg. Dallas 1, Texas

Neurology RAYMOND D. ADAMS, M.D.

Ophthalmology

Albert D. Ruedemann, M.D. Theo. E. Walsh, M.D. Detroit St. Louis

Internal Medicine WILLIAM B. CASTLE, M.D. E. P. PENDERGEASS, M.D. Milwaukee

**ROBING PROPERTY OF THE PROPERTY O

George C. Griffith, M.D. Los Angeles CYRUS C. STURGIS, M.D.
Ann Arbor

CORTES J. LUND, M.D.
New Orleans

Urology ELMER HESS, M.D. Erie

Ophthalmology

JAMES H. ALLEN, M.D. GEO. E. SHAMBAUGH, M.D. Chicago

Radiology

Pediatrics HORACE L. HODES M.D. New York City

Surgery O. T. CLAGETT, M.D. Rochester ARTHUR W. ALLEN, M.D.

Orthopedics

Surgery of the Hand Joseph H. Boyes, M.D. Los Angeles

GEORGE N. AAGAARD, M.D. Dean, Southwestern Medical School of the University of Texas, Dallas

J. P. Sanders, M.D. President, American Academy of General Practice

I, a member in good standing of the County Medical Society of the State of ... wish to attend the Spring Clinical Conference. Enclosed is my registration fee of \$20.00, which entitles me to all features of the meeting. .M.D. My practice is.... I am a graduate of; Year of Graduation...




The Emory John Brady Hospital 401 Southgate Road

A Private Hospital for Nervous and Mental Diseases

Situated in a beautiful valley two miles south of Colorado Springs, which is nationally known as a health center. New building for mild cases of Functional Neurosis, affording complete classification of patients. Home-like surroundings, scientific medical treatment and nursing care. Booklet and rates on application.

C. F. Rice, Superintendent, Colorado Springs, Colorado



GREATER BILE VOLUME



In biliary dysfunction, where copious flow of thin bile is the chief aim of therapy, the physician can place complete reliance upon DEHYDROCHOLIC ACID - Breon and SODIUM DEHYDROCHOLATE - Breon.

Because they are the least toxic of all bile acids and their salts, these products may be employed in full dosage to exert maximum therapeutic effect.

For most efficient drainage of the biliary tract these Breon preparations are indicated in chronic cholecystitis, noncalculous cholangitis, and other biliary disorders where no total obstruction is present.

In your management of biliary patients, specify "Breon" products for purity, pharmaceutical elegance, and uniform end results.

For Oral Administration Tablets - 0.25 Gm. each

For Parenteral Administration 20% w/v Solution



DEHYDROCHOLIC ACID - Breon

Bottles of 100, 500 and 1000 Tablets

SODIUM DEHYDROCHOLATE - Breon

5 cc. Ampuls - boxes of 6 and 25

Write Dept. MA 27M for literature.

George A. Breon & Company

Manufacturing Pharmaceutical Chemists NEW YORK 18, N. Y.

1450 BROADWAY



A Complete Production Service

ELECTROTYPES
MATRICES
STEREOTYPES
PRINTING
TYPOGRAPHY

Western Newspaper Union

Denver - - - - 1830 Curtis St.
New York - - - 310 East 45th St.
Chicago - - - 210 So. Desplaines St.

And 33 Other Cities



Juberculosis Abstracts

Issued Monthly by the National Tuberculosis
Association

Vol. XXV

JANUARY, 1952

Wa 1

CHEST X-RAYS ON ADMISSION PAY OFF

William Siegal, M.D., Robert E. Plunkett, M.D., and Herman E. Hilleboe, M.D., The Modern Hospital, July, 1951

Routine chest x-ray examination of patients admitted to general hospitals is a fruitful method of finding new cases of pulmonary tuberculosis. The patient, the hospital staft, and the community all reap benefits. The procedure yields greater returns in discovering unsuspected disease than mass x-rays of the general population or school groups.

In New York State, outside of New York City, there are 166 voluntary, nonprofit, and publicly supported general hospitals which annually admit over 650,000 patients. The plan prepared by the New York State Department of Health and supported by public funds was developed cooperatively through the Health Department and these hospitals. Policies and procedures were set up in 1946 and briefly are as follows:

Any nonprofit general hospital with an inpatient admission rate sufficiently large to provide 4,000 admission chest x-rays annually is eligible to borrow complete photofluorographic equipment for taking 4" x 5" or 70 mm. films. The hospital received 50 cents for each report of an admission x-ray film submitted to the local health department. The department recommends that hospitals install equipment as close to the admitting rooms as is practicable in order to maintain a high percentage of x-rays on admitted patients.

Hospitals whose admission rate is less than 4,000 patients annually may also participate in the program by using their own equipment. For this service, they receive one dollar for each x-ray report submitted. Of the 166 general hospitals, fifty-eight are eligible for loan of photofluorographic equipment and the remaining 108 can participate by using their own equipment. The fifty-eight hospitals eligible for loan of equipment represent only 37 per cent of all the hospitals, but account for 67 per cent of all the admissions.

Any participating general hospital, in applying to the State Health Department, agrees that it will:

1. Make every effort to x-ray the chest of all admitted patients, 15 years of age and over.

2. X-ray the chests of all employees not previously x-rayed and of all new employees.

WANTADS

GENERAL SURGEON-38, experienced, Boards and College, desires location, group preferred. Availale now. Write Box 3, Rocky Mountain Medical Journal.



H-O-W-D-Y

Reg. Trade Mark

BOB'S PLACE

A Bob Cat for Service

Be Western; Come Out to Cowtown— The Howdy Town. Your Drug Store Cowboy.

CONOCO PRODUCTS


300 So. Colorado Blvd.

Denver, Colo.

Save Infant Life with the

OXYGEN-AIR PRESSURE LOCK

(Bloxsom*)



An intermediate oxgen-air lock for the asphyxiated and premature infant between intra-uterine life and life immediately following delivery.

Infant mortalities are reduced

when the Oxygen-Air Pressure Lock is standard equipment in efficient hospitals.

Why not investigate the advantages of the Oxygen-Air Pressure Lock today?

*Allan Bloxsom, M.D.
The Journal of Pediatrics
Vol. 37 No. 3—Pages 311-319, Sept. 1950.

GEO. BERBERT & SONS, INC.

1524 Court Place DENVER 2, COLORADO 3. Make no charge to the patient for the initial x-ray examination and interpretation or for additional x-rays or services necessary to establish a diagnosis of tuberculosis.

4. Use the recommended diagnostic classification.

Submit an x-ray report for each patient and employee examined under this program to the local health official.

It was not expected that uniform procedures for the routine x-raying of admissions would be possible for all the hospitals. The following routine, however, was suggested and is being carried out with minor changes. Identifying information is entered on a special report form at the time of admission for every patient 15 years of age or over. If possible, he is x-rayed, usually without disrobing, before being taken to his room. If he is too ill to be r-xayed on admission, this is done as soon as his physical condition permits. The admission films, 4" x 5", 70 mm. or 14" x 17", are processed and interpreted within twenty-four hours and the diagnoses, if negative or nontuberculous, are checked on a special report form.

If the admission film shows definite or suspected tuberculosis, additional chest x-ray and other examinations are made for diagnostic and clinical evaluation. The diagnosis is entered on the admission x-ray report. Completed admission x-ray reports are sent frequently to the health officer. In addition, the hospital also furnishes the health officer with a monthly bill for the admission chest x-ray reports submitted to him. If active tuberculosis is found, the hospital then makes an official

case report.

It is important that the hospitals use the same classification of disease, especially as it relates to tuberculosis, in reporting the results of these x-rays. The admission small or large film diagnosis is not considered the final diagnosis or determination of activity. Nevertheless, a tentative diagnosis is necessary in case patients do not remain in the hospital long enough for further detailed study when it is indicated. The health officer should know what persons with possible tuberculosis return to the community from the hospital. A tentative diagnosis, therefore, is made on all films. If the tentative diagnosis is definite tuberculosis, an estimate of clinical status is also made. If probably active, the extent of the disease is also noted. Films which indicate pleural effusion otherwise unexplained are considered to be probably active tuberculosis.

The follow-up of cases of definite and suspected tuberculosis found by the hospital is the health officer's responsibility. The admission x-ray reports sent to the health officer are a check on the number billed by the hospital for reimbursement; they are used also for detailed monthly reports which are sent to the State Department of Health. The health officer maintains a separate file of positive x-ray reports and arranges for follow-up examinations. These include provision for diagnostic and clinical determination for each person reported and adequate medical care. For each report of a definite or suspected case of tuberculosis, the health officer submits to the department at the end of six months, a summary of what has happened to the person during the interval.

An analysis of the initial chest x-ray examinations of adults admitted to the general hospitals participating in this program from May, 1947, to January, 1950, shows

A total of 195,751 patients, 15 years of age and over, had chest x-rays taken on admission to forty-one general hospitals. The largest number of patients examined (48 per cent of the total) was in the age group 15 to 34. Females outnumbered males two to one; the ratio of females to males was in excess of four to one between the ages 15 to 34. From the initial hospital x-ray interpretations, 3,976 or 20.3 for every 1,000 patients x-rayed, were tentatively diagnosed definite or suspected tuberculosis. Of these 1,005 or 5.1 per 1,000 x-rayed, were considered to have probably active pulmonary tuberculosis. For all ages the prevalence of probably active pulmonary tuberculosis was three times as great among males as among females, the highest prevalence being in males 45 years of age and over.

The distribution of the probably active cases by stage of disease was: minimal 47 per cent, moderately advanced 35 per cent, and far advanced 18 per cent. Fewer minimal and more advanced cases are found in general hospital patients than in community surveys.

general hospital patients than in community surveys.

Of 126,190 admission chest x-rays during the period between January, 1948, and June, 1949, inclusive, 2,642 showed evidence of definite or suspected tuberculosis, of which 2,145 had not been previously reported and were considered new cases. Of these, 71.7 per cent received follow-up examinations within six months of the initial hospital diagnosis.

On the basis of the number diagnosed definitely active,

NEWTON OPTICAL COMPANY

GUILD OPTICIANS

309-16th Street

Phone KEystone 0806

Catering to Medical Profession Patronage

Denver

We value the business of the many doctors we serve.


MERCHANTS OFFICE FURNITURE COMPANY

1511 Arapahoe Street

AComa 2559

Denver, Colorado

Don't miss important telephone calls .



Let us act as your secretary while you are away, day or night: our kindly voice conscientiously tends your telephone business, accurately reports to you when you return.

Telephone ANSWERING Service CALL Alpine 1414

for normal infant development

Clinical experience with thousands of infants demonstrates impressively the valuable role of Bremil in infant nutrition.


Bremil is a completely modified milk in which nutritionally essential elements of cow's milk have been adjusted in order to supply the nutritional requirements of infants deprived of human milk. It can be used with confidence either as part or all of the food supplied to the normal healthy infant.

Bremil conforms to the fatty acid and amino acid patterns of human milk. Bremil is a completely modified milk in which the calcium-phosphorus ratio (guaranteed minimum 11/2:1) is adjusted to the pattern of human milk, thus helping to prevent tetanic symptoms in newborns. 1,2

Bremil supplies the same carbohydrate as breast milk, lactose.3

Bremil's vitamin adjustments for standards of infant nutrition,4 its human-milk size particle curd, miscibility and palatability are additional reasons for its choice in infant feeding. Bremil approximates the nutritional role of the mother.





2 Nesbit, H. T.: Texas State J. M. 38:551, 1943

3 Bull. National Research Council No. 119 Jan. 1950

4 Recommended Daily Dietary Allowances, Revised 1948, Food and Nutrition Board, National Research Council

Complete data and Bremil samples are available to you

Prescription Products Division

The Borden Company



palatable,

easy

prepare

350 Madison Avenue, New York 17

e e e a ror nof hix

of in

al 48 4. of en

er-

re er-

ve

ng

ge id-

nt.

in

42 of

ere

IAL

activity undetermined and suspicious for tuberculosis after follow-up, it is estimated that, if adequate follow-up had been possible for all the 2,145 new cases of definite or suspected tuberculosis, a ratio of 2.5 active cases of tuberculosis would be found for every 1,000 patients x-rayed.

Vol. XXV

FEBRUARY, 1952

TREATMENT OF MINIMAL TUBERCULOSIS Iulia Iones, M.D., The NTA Bulletin, September, 1951.

All pulmonary tuberculosis is minimal in its early stages and progression may be avoided by effective treatment. Minimal disease is usually unaccompanied by symptomatic illness so is frequently demonstrated only by roentgenogram. The discovery of disease in its early and curable forms is one of the purposes of case-finding programs. While the wisdom of these programs is established, they fall short of complete accomplishment unless subsequent treatment is effective. Once minimal disease has been identified, its progression must be accepted as failure.

Because the lesions are small and the patient has few, if any, symptoms, one is tempted to attach less sig-nificance to minimal disease than to the advanced disease. Rather than to reassure himself and the patient by thinking of "just a little spot on the lung," the phy-sician needs to consider the lesion as the focal area from which disabling disease may occur and to take advantage of the opportunity for simple, less costly, and more effective treatment.

Individuals vary in their ability to heal tuberculosis infection. The majority of those first infected with the disease remain well and infection is indicated only by reaction to tuberculin. Small areas of calcification may appear eventually in lung or lymph nodes. In others irreversible damage occurs, and necrotic tissues may

liquefy and slough into nearby bronchi and healthy lung. While the patient may be without symptoms, the x-ray reveals small shadows of pneumonic disease and a 'minimal" lesion is diagnosed. The patient may then develop sufficient resistance to prevent further extension. In this case, he either remains well or may harbor areas of chronic infection which undergo evolution after considerable lapse of time.

Except by hindsight, it is not possible to distinguish between the individual who can control his minimal lesion without treatment and the patient in whom progressive disease may occur. This often proves costly for the patient and the community. Since some undetected lesions are controlled without treatment, residual shadows may later be encountered in routine x-ray examinations. For this reason various factors including the pathologic age and character of lesions discovered in asymptomatic individuals need consideration before treatment

Previous x-ray examinations may demonstrate that the lesion is newly acquired and must be assumed active and unstable. Symptoms or the presence of tubercule bacilli in sputum or gastric contents may indicate activity. In adolescents and young adults, most minimal disease is recently acquired and quite unstable. While new disease may be acquired throughout life, lesions occurring in older persons may represent old unidentified disease needing only periodic examination. Lesions must be subjected to clinical scrutiny establishing their duration and potentialities. From these studies will emerge those patients with early, minimal disease. Effective treatment of this group constitutes the major problem in dealing with minimal disease.

Early lesions are small areas of tuberculous bronchopneumonia which may resolve completely, leaving essentially normal lung tissue. On the other hand, the tissues within this area may be destroyed, leaving cavities from

LIVERMORE SANITARIUM



- The Hydropathic Department devoted to the treatment of general diseases, excluding surgical and acute infectious cases. Special attention given functional and organic nervous diseases. A well equipped clinical laboratory and modern X-ray Department are in use for diagnosis.
- · The Cottage Department (for mental patients) has its own facilities for hydropathic and other treatments. It consists of small cottages with homelike surroundings, permitting the segregation of patients in accordance with the type of psychosis. Also bungalows for individual patients, offering the highest class of accommodations with privacy and comfort.

GENERAL FEATURES

Climatic advantages not excelled in United States. Beautiful grounds and attractive surrounding country.
 Indoor and outdoor gymnastics under the charge of an athletic director. An excellent Occupational Department.
 A resident medical staff. A large and well-trained nursing staff so that each patient is given careful individual attention.

Information and circulars upon request. Address: O. B. JENSEN, M.D. Superintendent and Medical Director LIVERMORE, CALIFORNIA Telephone 313

CITY OFFICES:

SAN FRANCISCO 450 Sutter Street GArfield 1-5040

OAKLAND 1624 Franklin Street GLencourt 1-5988





EFFECTIVE and OPERATING FOR

Members of your State MEDICAL PROFESSION

DESIGNED EXCLUSIVELY By CONTINENTAL CASUALTY COMPANY

One of the Oldest and Largest Stock Casualty Companies in the World

A Few of the Many Outstanding Features

- ★ Monthly benefits for both accident and sickness payable FROM FIRST DAY Minimum claim 7 days.
 ★ Pays monthly benefits for disability resulting from "accidental bodily injury": —the term "accidental means" is NOT
- Pays monthly accident benefits regard-less of whether disability commences within a certain specified time.
- * No automatic termination age.

- ★ The premium does not increase as you grow older.
- House confinement is never required to collect full benefits.
- Indemnities cannot be prorated because of other insurance or for performing duties of a more hazardous occupation.
- * Policies cannot be restricted by rider to exclude any disease or injury originating after date of issue.

Renewal is guaranteed to individual active members of the profession regardless of age, so long as the premiums are paid in accordance with the terms of the contract and the plan continues in effect for the members in your designated territory.

The Continental Casualty Company pioneered in the writing of Professional Association Insurance plans. The first such plan, written in 1923, is still in effect and the Company has never cancelled or declined to renew a Professional Plan because of unfavorable loss experience.

NOW available to all eligible members in active practice and under 69 years of age.

Plans A or B available to female members under age 65.

SICKNESS BENEFITS	PLAN AA	PLAN A	PLAN B	
PAYS A Monthly Indemnity for total disable during first year, whether house confor not,	ined	\$ 200.	\$ 100.	
PAYS Total Monthly Indemnity First 2 Years	5400.	3600.	1800.	
ACCIDENT BENEFITS				
PAYS A Monthly Indemnity for total disab during first yearFrom First		200.	100.	
PAYS Total Monthly Indemnity First 2 Years	5400.	3600.	1800.	
PAYS Monthly Indemnity for partial disabup to 13 weeksFrom First		80.	40.	

The Cost is Low

	PLAN AA	PLAN A	PLAN B
Annual	\$138.00	\$92.00	\$46.00
Semi-Annual	69.50	46.50	23.50

ADD \$5.00 TO FIRST PREMIUM ONLY

OPTIONAL BENEFITS

PLAN AA PLAN A PLAN B

\$ 150.

Monthly Hospital Indemnity may be added up to for \$6.00 per \$100 Hospital Benefits payable for \$ 450. \$ 300. for \$6.00 per \$100 Hospital One to 90 days — Each disability.

Accidental Death and Dismemberment Indemnity up to.. 2500. 7500. 5000. Added for \$2.00 per \$1000.

CONTINENTAL CASUALTY CO.

PROFESSIONAL DEPARTMENT, Intermediate Division 30 EAST ADAMS STREET - CHICAGO 3, ILLINOIS

AL

which dissemination may occur. Even in the smallest lesion demonstrable by x-ray, some areas have undergone destructive changes. The outcome of any case depends upon the extent and character of the disease, mdividual factors of resistance, and the manner in which the latter are influenced by treatment.

The therapeutic program is developed from the following considerations. Since the minimal lesion represents recent extension of disease from microscopic foci of greater duration, it must be assumed that the patient, at this stage, has inadequate resistance to control his disease. Treatment must be directed toward increasing resistance. Much of the lesion may be reversible if the lesion has been discovered soon after it has developed. Immediate treatment is urgent before further evolution produces less reversible lesions. If further extensions of disease take place, each new lesion has potentialities for breakdown and further dissemination. The presence of small necrotic foci must be assumed in all minimal lesions and their extent limits the effectiveness of cure.

Rest is the foundation of the therapeutic program. Experience indicates that rest favors development of resistance, thus enabling the tissues to suppress activity of the tubercle bacillus, remove products of inflammation, and to control areas more permanently damaged. Spreading disease occurs less often when patients are in bed. Bronchial secretions are decreased during bed rest, and this factor probably plays an important part in decreasing the hazard of dissemination through the bronchi.

Bed rest is most effective during the carly period of treatment when the lesions are reversible and most unstable. For this reason, it is advocated that patients with early lesions be put to bed immediately upon identification of their lesions. Often this is difficult since the patient feels well. Compromises which permit him to continue his normal activity while the lesion is observed, may jeopardize his future health and happiness.

observed, may jeopardize his future health and happiness. It is difficult for a asymptomatic individual to make the transition from an active life to complete rest. Given thorough understanding of his problem and the odds at stake and given day to day assistance in meeting the aggravations of inactivity, the usual patient is less unhappy from his treatment than from a set-back of progressive disease. An intelligent individual is able to accept the depressing aspects of tuberculosis infection and inactivity. Recognition of individual problems is necessary and special assistance may be needed.

It seems wise to continue bed rest until stability of the lesion can be assumed. This implies absence of constitutional symptoms and an unchanging lesion by roentgenogram. Clearing of reversible elements occurs usually in from four to six months. Subsequent change by x-ray may be slight and quite slow. From this point treatment is directed toward control of more permanently damaged areas whose presence must be assumed. The time necessary depends on the patient's clinical course, personal situation, and anticipated demands of his normal activities. Resumption of activity must be gradual since bed rest is deconditioning.

In some cases, the administration of streptomycin and para-aminosalicylic acid may be wise. But bacterial resistance may develop and, since the minimal lesion is potentially the advanced lesion, an effort must be made to conserve this temporary support for urgent needs.

Most patients recover permanently from minimal disease if rest is adequate. A few develop more chronic disease which continues to threaten health, and in this group it may occasionally be necessary to add collapse or other surgical therapy.

Effective treatment of minimal tuberculosis must be prompt and thorough. The patient must be thoughtfully taught about his disease if full cooperation is to be achieved. Most patients recover completely and resume their previous activities, but needs for vocational retraining must be visualized. Regular medical supervision wisely continues after recovery and resumption of normal living.

We Appreciate the Patronage of the Members of the Medical Profession

CAPITOL SANDWICH CO.

Established 1921

Sandwiches on Sale at the Better Drug Stores of Denver

KEystone 2694 or EAst 4707

Denver

Colorado

Established 1894

Paul Weiss

OPTICIAN 1620 Arapahoe Street Denver, Colo. 50 Years of Ethical Prescription Service to the Doctors of Cheyenne

*

ROEDEL'S PRESCRIPTION DRUG STORE

RESCRIPTION DRUG STOR

CHEYENNE, WYOMING

Bonita Pharmacy

(Established 1921)

Prescription Pharmacists

6th Avenue at St. Paul Street "RIGHT-A-WAY" SERVICE

GERALD P. MOORE, Manager Phone FRemont 2797

PRESCRIPTION BLANKS

ONLY 5725 on orders for 5,000 Smaller Quantities

3,000 @ \$2.85

per 1,000

1,000 for \$3.45

3

n n is

of

'nt

d

al of

e

in al

le

al

ic

is

OI

be

lly

be

ne

re-

vi of

TAL

Printed by experienced craftsmen on Pur-White Sulphite Bond paper, in the distinctive Modern Futura type face. The 4" x 51/2" blanks come in pads of 100, shipped within 1 week after receipt of your order. The product delivered, is belitting the dignity of your profession.

FREE DELIVERY ANYWHERE IN THE U.S.A. Send Check or Money Order (No C.O.D. please)

GENUINE LEATHER COVER

FOR YOUR RX BLANKS

WITH EVERY ORDER FOR 5,000 (FOR A LIMITED TIME ONLY)

Your initials will be stamped in gold on the Leather Rx Blanks Cover.

MONEY REFUNDED IF NOT COMPLETELY SATISFIED Write for Free Folder on other printed items

Fine Printers Since 1897

CO. 27 W 24th St., NY 10, NY

DEEP ROCK Artesian Water

Famous for over 52 years as Denver's finest and purest drinking water.

- · Endowed by Nature with the ideal amount of fluorine, 1.3 parts per million
- Contains no added chemicals
- Recommended by Doctors for baby formulas, stomach and kidney disorders

DEEP ROCK Distilled Water

- Scientific distilling process removes all minerals
- · Aerated, to remove flat taste of other distilled waters
- Recommended by Doctors for baby formulas, allergies, prescriptions and sterilizing

Order New At Your Pharmacists or call TAbor 5121

DEEP ROCK WATER CO. Denver, Colorado 614 27th Street

YORK PHARMACY

Denver's Finest Prescription Store

J. GLEN MATSON, Owner Free Delivery Phone FR. 8837

2300 East Colfax Avenue at York Street **Almay Cosmetics**

NURSES' OFFICIAL REGISTRY

Endorsed by District No. 2 Colorado State Nurses' Association American Nurses' Association

REGISTERED NURSES PRACTICAL NURSES

Nursing Service for All Community Needs

KEystone 0168 **Argonaut Hotel**

Colfax and Grant, Denver

RADIUM AND RADIUM D + E

(including Radium Applicators)

For All Medical Purposes Est. 1919

QUINCY X-RAY & RADIUM LABORATORIES

(Owned and Directed by a Physician-Radiologist)

HAROLD SWANBERG, B.S., M.D., Director

W. C. U. Bldg.

Quincy, Illinois

Denver's Fireproof

COLBURN HOTEL

D. B. Cerise is the genial Host and Manager

- CONVENIENT Located only a ten-minute walk from the heart of the city.
- PLEASANT Away from above the noise and rush of downtown Denver.
- EXCELLENT FOOD Dining that has satisfied the demanding tastes of all partons.
 Visit Our New Cocktail Louage.

TENTH AVE. at GRANT ST.

Phone MAin 6261

Denver, Colo.

WALTERS DRUG STORE 801 COLORADO BLVD.

Denver, Colorado



Telephone FRemont 5391

Welcome to

Aylard's Crestmoor Drug

3rd and Hudson

Prescriptions — Biologicals — Chemicals

Drugs and Sundries

Cosmetics

Soda Fountain

Phone FLorida 2326

Denver, Colo.

GABRIEL'S RESTAURANT 240

MISS M. E. GABRIEL, Prop.

SERVING TRADITIONALLY GOOD FOOD AT MODERATE PRICES

HOURS: 11:00 A.M.—2:00 P.M. 4:30—7:30 P.M. SUNDAYS: 12 Noon to 7:00 P.M. Closed Wednesdays

240 Broadway

Denver, Colo.

SPruce 2182

We Cater to the Medical Profession

CASCADE LAUNDRY

10 Per Cent Discount If You Bring Your
Laundry

HAND DRY CLEANING

"Deserving of Your Patronage"

618 East 16th Ave., Denver TAbor 6379

Charge Accounts Inviteo



hen it is impossible to take your product to the customer, or have him come to your establishment, you will find it both impressive and profitable to show your product by picture.



Ready Now!

4th Edition STANDARD NOMENCLATURE of DISEASES and OPERATIONS

Edited by Richard J. Plunkett, M.D., and Adaline C. Hayden, R.R.L.

1,034 pages. Illustrated. \$8.

Stacey's carries the medical and technical books of all publishers. You are cordially invited to phone AComa 3411, drop in and browse, or write for any of your book requirements.

Stacey- TECHNICAL BOOK CO. 1814 STOUT STREET DENVER 1. COLORADO

A Western Institution

RELIABLE DRUGGISTS

PATRONIZE DENVER'S INDEPENDENT DRUGGISTS

North Denver's Largest Rx Stock CALL GLendale 3643-3644

Ask for Rx Department

Qualified Registered Pharmacists
Answer the Phone

We Deliver to Any Part of North Denver Any Time

> WOODMAN PHARMACY 4400 Tennyson Street

24 Years in the Heart of North Denver

GUIDO SHUMAKE DRUGS

PRESCRIPTIONS ACCURATELY
COMPOUNDED

Free Delivery Service

West 38th Ave. and Clay Denver, Colo.

Phone GLendale 1073

For Accurate Prescriptions-

For prompt delivery thruout the area-

Phone: BElmont 3-4621

g

Kincaid's Pharmacy

7024 W. Colfax Ave., Lakewood, Colo.

We Recommend

BONNIE BRAE DRUG COMPANY

Alfred C. Anderson, Owner and Manager

Prescriptions Accurately Compounded

Drugs . . . Sundries

Complete Line of Cosmetics

FREE DELIVERY

763 South University Boulevard

Phone RAce 2874 - Denver, Calorade

WE RECOMMEND

Whittaker's Pharmacy

"The Friendly Store"

PRESCRIPTION SPECIALISTS

West 32nd and Perry, Denver, Colo.

Phone GLendale 2401

HYDE PHARMACY

ACCURATE PRESCRIPTIONS

Chas. W. Hyde, Prop.

Rocky Mountain Distriutors for Sherman Biologicals and Pharmaceuticals Almay Non Allergic Cosmetics

Prompt Free Delivery

KE. 4811

WA 4566

1400 East 18th Avenue at Humobldt

WE RECOMMEND

LAKEWOOD PHARMACY

R. W. Holtgren, Prop.

PRESCRIPTION SPECIALISTS

West Colfax at Wadsworth

Lakewood

Colorado

Phone BElmont 3-6531

L K PROFESSIONAL

Phone Aurora 1900 or Dial FLorida 1864 9350 East Colfax Avenua

Specializing in Prescriptions

Free Delivery in Aurora Area

Almay Hypoallergic

Cosmetics

Flaherty Surgical Supports

Lou and Ken Suher

WHEATRIDGE FARM DAIRY

"Pride of the West"

Ice Cream for All Occasions

COMPLETE LINE OF GRADE "A"
DAIRY PRODUCTS

HOMOGENIZED MILK

8000 West 44th Ave.

GL. 1719

ARVADA 220

HAVEN PHARMACY

J. L. Panek, Jr., Prop.

PRESCRIPTION DRUG STORE

DRUGS AND SUNDRIES

29th and Irving St. Phone GLandale 5191

We Make Free Prescription Deliveries



ARTIFICIAL EYES

Serving the doctor and his patient with the finest in natural appearing artificial eyes since 1906. Plastic eyes made to order. Largest selection of glass and plastic eyes in America. Specialists in building eyes for all types of implants. Write or phone for full details.

DENVER OPTIC CO., 330 University Bldg., 910 16th St., Denver 2. Ph. MA. 5638

Oculist Prescription Service Exclusively

SHADFORD-FLETCHER OPTICAL CO.

Dispensing Opticians
228 16th Street, Denver, Colo. AComa 2611
3705 East Colfax (Medical Center Building). FLorida 0202

Stodghill's Imperial Pharmacy

Prescriptions Exclusively

For your prescriptions we stock a complete line of ALMAY—non-allergic—cosmetics.

Five Pharmacists

319 16th St.

TAbor 4231

Denver, Colo.



WINNING HEALTH
in the
Pikes Peak Region

COLORADO SPRINGS

Inquiries Solicited

GLOCKNER PENROSE HOSPITAL

Sisters of Charity
HOME OF MODERN SANATORIA

Index to Advertisers

Pag	ge	Page	Page
Abbott Laboratories 14	43	Fairfax Sanitarium 100	Republic BuildingCover 3
American Medical and		Florida Citrus Commission 108	RMS Printers 159
Dental Association	94		Roedel's Prescription
Averst, McKenna &		Gabriel Restaurant 160	Drug 158
Harrison 13	33	Glockner Penrose Hospital 162	
Aylard's Crestmoor Drug. 1	60		St. Anthony Hospital 148
		Haven Pharmacy 162	Schering Corporation 93
Baker Laboratories 19 Berbert, George &	47	Hyde's Pharmacy 161	Searle, G. D., & Co
Sons, Inc 1		Kendrick-Bellamy Co 92	Optical Co 162
Bob's Place 1		Kincaid's Pharmacy 161	Shirley-Savoy Hotel 148
Bonita Pharmacy 1			Shumake Drug, Guido 161
Bonnie-Brae Drug 1		V V Destauries	
Borden Company		L K Professional Pharmacists	Stacey-Technical Books
Breon, George A. & Co 1			Co., Inc
Brown School 1	42	Lakewood Pharmacy 161	Stapleton, H. C., Drug Co. 148
		Lederle Laboratories 135	Stodghill's Imperial
Cambridge Dairy	98	Lilly, Eli & CoCover I	Pharmacy 162
Camel Cigarettes 1		Lilly, Eli & Co109-110	Squibb, E. R., & Sons106-107
Capital Chevrolet 1		Livermore Sanitarium 156	
Capitol Sandwich Co 1			Technical Equipment
Carlson-Frink		Mead Johnson &	Corp 145
Cascade Laundry 1		CoCover IV	Telephone Answering
Children's Hosp. Assn 1 City Park Dairy 1		Merchants Office	Service 154
Colburn Hotel 1		Furniture Co 154	Thornton, George R 92
Colo. Artificial Limb	100	Merck & Company 105	and the same of th
Cp., Inc	148	Mercy Hospital	
Colvin Medical Books 1			United States Brewing
Continental Casualty Co 1	157	Morning Milk 97	Industry140
Cook County Graduate	+		Upjohn Co 95
School of Medicine 1	144	Newton Optical Co 154	
		Nurses Official Registry 159	Walters Drug Store 160
Dallas Southern			Wander Company 137
	150	Park Floral Company 102	Wantads 152
Davis Bros. Drug Co		Parke, Davis &	Weiss, Paul 158
Deep Rock Water		CoCover II-91	Western Newspaper Union 152
Denver Optic Company	162	Peters, Winter &	Wheatridge Farm Dairy 162
Denver Towel Supply Co	146	Christensen, Inc 136	Whittaker's Pharmacy 161
Dorr Optical Co	104		Winthrop-Stearns, Inc 99
		Physicians Casualty Assn 144	
Earnest Drug Company	149	Professional Pharmacy 102	
Ehret Engraving Co			Wyeth, Inc 10
Emory, John Brady	100	Quincy X-Ray and Radium	
Hospital, The	150		York Pharmacy 159
ALUDPIUM, AME	200	Landiatories	



Woodcroft Hospital-Pueblo, Colorado

A private hospital for the scientific treatment of neuro-psychiatric disorders, including alcoholism and drug addiction. Beautiful landscaping and home-like surroundings afford a restful atmosphere. Acommodations vary from single rooms with or without bath to rooms en suite, allowing for segregation of guests.

Detailed information furnished on request. Karl J. Waggener, M.D.



THE CHILDREN'S HOSPITAL ASSOCIATION of DENVER

NON-SECTARIAN-NON-PROFIT

Providing medicinal and surgical aid to sick and crippled children of the Rocky

Mountain Region from Birth to Maturity

Every modern scientific aid available to the physicians and surgeons of Colorado and Wyoming

Approved by the American Medical Association and the American College of Surgeons Full Three-Year Nurses' Training Course

The Republic Building

DENVER'S OUTSTANDING
MEDICAL CENTER



Designed for the exclusive use of the Medical and Dental Professions, the Republic Building is the largest medical building in the Rocky Mountain region, serving families from throughout Colorado and the surrounding states.

Barres ons

to build a sound infant formula

Openerous Protein Fat Content
Appropriate Fat Conbohydrate
Adequate Conbohydrate

LACTUM has these three dimensions

...for Lactum is an evaporated whole milk and Dextri-Maltose® formula.

Its proportions are those of milk and Dextri-Maltose formulas used successfully in infant feeding for forty years.

And its caloric distribution (16% protein, 34% fat, 50% carbohydrate) is based on authoritative pediatric recommendations.



and Lactum has a 4th dimension

...time-saving convenience
Lactum feedings are prepared
simply by adding water,
A 1:1 dilution provides
20 calories per fluid ounce,

For Premature and full Torm
Infants with Low Fat Tolerance
Infants with Low Fat Tolerance

DALACTEM. Mead's evaporated to the formula, offers the same of the formula, offers the policy formula, offers the f

MI



MEAD JOHNSON & CO. EVANSVILLE 21, IND., U.S. A.

